Berkeley Faculty Roundtable on Environmental Services in Rangeland Production Systems


Presentation and Discussion Notes from the Third Roundtable: May 28, 2009


PATRICK GONZALEZ

MONITORING CARBON IN SAVANNAS AND WOODLANDS WITH FIELD INVENTORIES AND REMOTE SENSING

Berkeley Faculty Roundtable on Environmental Services in Rangeland Production Systems

Part I:
Powerpoint Presentation by Patrick Gonzalez





Presentation Outline

- I. Carbon estimation and monitoring process
- 2. Example from the African Sahel
- 3. Baseline estimation and sequestration projection



Presentation Outline

- I. Carbon estimation and monitoring process
- 2. Example from the African Sahel
- 3. Baseline estimation and sequestration projection



Greenhouse Gas Categories and Methods

Greenhouse Gases^A Carbon Pools^A Land Use^A

Aboveground Biomass^D

Forest land CO₂ Aboveground biomass

Mature trees

CH₄ Cropland

Deadwood

Understory trees

Grassland N₂O

Litter

Shrubs

Wetland **HFCs** Harvested wood products Grasses

Settlements PFCs

Belowground biomass

SF₆ Other land

Soil organic carbon^B

Carbon Methods^A

Uncertainty Methods^A

Gain-Loss (fluxes)

Simple error propagation

Stock difference

Monte Carlo

Sources

- IPCC 2006. National Greenhouse Gas Inventory Guidelines
- W. Silver, March 6, 2009, Rangeland Roundtable
- D. Baldocchi, March 20, 2009, Rangeland Roundtable
- P. Gonzalez, May 28, 2009, Rangeland Roundtable



Center for Forestry, University of California, Berkeley

Research Objective

Estimate additional carbon sequestration ($\Delta C_{\text{additional}}$) from natural resource management practices through estimates of carbon stocks (C_t) or fluxes in a defined area at specific times (t), compared to baseline changes ($\Delta C_{\text{baseline}}$)

$$\Delta C_{\text{additional}} = \frac{C_2 - C_1}{t_2 - t_1} - \Delta C_{\text{baseline}}$$

t_1	t_2	$\Delta \mathcal{C}_{ ext{additional}}$	$\Delta C_{baseline}$
30 years ago	this year	observed trend	0
last year	this year	monitoring	0
this year	30 years from now	projection	projection



Carbon Estimation and Monitoring Process


- Delineate area (landscape or project)
- 2. Estimate current carbon stock
- 3. Monitor stocks over time by repeating methods
- 4. Estimate baseline sequestration
- 5. Project future sequestration





Presentation Outline


- I. Carbon estimation and monitoring process
- 2. Example from the African Sahel
- 3. Baseline estimation and sequestration projection












Approximate Scale: 1:10,000,000


Africa Land Cover


Bartholomé and Belward 2005. Int.'l J. Remote Sensing




Research Area, Northwest Senegal, 7600 km²

Gonzalez 2001. Climate Research









Forest Species Loss

ca. 1945 64 ± 2 species

1993 43 ± 2 species

change -33% ± 5%

significance p < 0.001


sample $n = 135 \times 4 \text{ km}^2$


Sahel, Sudan, Guinean zones shift SW 25-30 km


Gonzalez 2001. Climate Research


Tree Dieback, Northwest Senegal, 1954-1989,

Trees of height > 3 m











Normalized Difference Vegetation Index (NDVI)




Tucker 1979. Remote Sensing of Environment

Aboveground biomass vs. integrated NDVI



Tucker et al. 1985. Remote Sensing of Environment



Other Estimation Parameters

- I. Carbon fraction of biomass
- 2. Root:Shoot ratios
- 3. Dead tree fraction of live tree
- 4. Coarse woody debris relationships
- 5. Litter relationships
- 6. Wildfire relationships






Senegal Sahel Carbon Estimate					
	Growth (Mg ha ⁻¹ y ⁻¹)		Gross rate (y-1)		
Total carbon	3.3	41	0.08		
Total vegetation					
herbaceous	2.9	3.2	0.91		
ligneous	0.2	12	0.02		
total	3.1	15	0.20		
Aboveground biomass and deadwood herbaceous 1.2 1.3 0.91					
ligneous	0.2	7.3	0.03		
total	1.4	8.5	0.16		
Belowground bior herbaceous ligneous total	nass 1.7 0.03 1.8	1.9 4.8 6.7	0.91 0.01 0.26		
Soil Woomer et al. 2004	0.2	25	0.01		
Uncertainty (SD) by simple error propagation ±30%					

IKONOS sites, Northwest Senegal, 200 km²

P. Gonzalez, C.J. Tucker, H. Sy, research in progress





IKONOS Njóobéen Mbataar, Senegal February 17, 2002

Image Space Imaging, Inc., Analysis P. Gonzalez





IKONOS Fété Olé, Senegal February 22, 2002

Image Space Imaging, Inc., Analysis P. Gonzalez







IKONOS Wolum, Mauritanie February 22, 2002

Image Space Imaging, Inc., Analysis P. Gonzalez






Njageen, Senegal, 1954, aerial photograph, IGN




Njageen, Senegal, 1989, aerial photograph, JICA




Njageen, Senegal, 2002, IKONOS satellite image






Tree Crown Estimates

Crown delineation



Aboveground biomass



In(biomass) =
$$18.613*((1-e^{-0.0757*diameter})^{0.2963})$$

 $n = 331, r^2 = 0.73$

P. Gonzalez, G.P.Asner, J.J. Battles, M.A. Lefsky, K.M. Waring, M. Palace, manuscript submitted



Monitoring Carbon in Savannas and Woodlands with Field Inventories and Remote Sensing

Presentation Outline

- I. Carbon estimation and monitoring process
- 2. Example from the African Sahel
- 3. Baseline estimation and sequestration projection



Principles of carbon sequestration project activities

Additionality

Property by which removal or sequestration of greenhouse gases results only from the direct action of a project activity, not from other activities already planned or funded

Baseline


Changes in greenhouse gas emissions and removals that would have occurred in the absence of the project activity


Leakage


Unintended displacement of greenhouse gas emissions to areas outside of the project boundary by actions attributable to the project activity



Baseline and sequestration projections for proposed forest project in Selva Central, Peru








P. Gonzalez, B. Kroll, C. Vargas, 2006, Report to U.S. Department of Energy







Part II: Discussion

Notes compiled and synthesized by Kayje Booker, Roundtable GSR

I. Issues and questions in response to Patrick's presentation:

- <u>Vegetation Indexes</u>: There are many indexes for measuring and monitoring vegetation. One drawback of NDVI is that it can saturate. Do you recommend NDVI?
 - NDVI works well for California rangelands because, although it can saturate if there are multiple stories, that situation is rare in rangelands.
 - EVI is another vegetation index using MODUS, but the advantage of NDVI is that there is data going back to 1980.
- Calculations of belowground biomass: Because the root to shoot ratio of trees and grasses are different, with grasses having relatively more of their biomass belowground, shouldn't the estimation of belowground biomass reflect that? Couldn't you turn the graph on slide 22 (Aboveground biomass vs. integrated NDVI) to a graph of total biomass by adding a line for belowground biomass that would be greater and have a steeper slope on the left hand side of the graph, where aboveground biomass is lower (because there most aboveground biomass would be grasses) then become flatter and lower at higher end of aboveground biomass (because there most aboveground biomass would be trees)? Has anyone validated that prediction?
 - It seems that that would be what you would expect and that such a regression and prediction would be possible. However, it has not been validated (as far as Patrick knows), and he advised caution on drawing too many conclusions from the regression.
- <u>Photo resolution</u>: What is the resolution on the 1954 aerial photos? Is it adequate for comparison with modern photos?
 - Resolution is limited on the older aerial photos. It improved greatly from 1954-1989, and IKONOS even better. The improvement of the resolution normally would create a problem in comparing across years because you can detect more trees now. In this case, however, it may be less of a problem, because Patrick has detected a trend of decreasing cover. If he is seeing decreasing tree cover even with better resolution, that finding is actually more robust. Still, he is working on standardizing between the different methods within the next two months.
- Species Richness: How do you detect species richness changes?
 - Field work with direct observation and interviews
 - The Wooloff have a profound knowledge of natural history. He did interviews with village elders and found less than 10% error in their recollections and recorded observations, so the interviews are quite accurate.

II. Broader discussion

Patrick started off the discussion by posing the following question: Given the low rate of increase of carbon in rangelands, is it worth it to pursue this option or would the money be better used in other carbon mitigation activities? Discussion centered around the following three broad themes: measurement and monitoring of carbon, forest regeneration, and managing for multiple uses including livestock grazing and carbon sequestration.

A. Measurement and monitoring of carbon

- Given the low rate at which carbon is sequestered in rangelands and the extreme heterogeneity of rangelands, isn't this whole discussion moot because the cost of monitoring carbon in rangelands outstrips the value?
- Depends on scale. If you could scale up to whole state of California, instead of
 monitoring plot by plot, and use remote sensing, rangelands as a whole in California
 could break even. For local sites, even in high biomass sites in tropical forest, it is hard
 to make money off of carbon because of management and monitoring costs. The
 smaller the project, the worse the money works out.
- As far as tools for that monitoring, Patrick thought that Landsat would be a good tool but that MODUS, at the state level, would probably not be adequate. It depends on the policy the level of allowable uncertainty dictates the resolution required.
- California Climate Action Registry (CCAR) has been developing a forest protocol, and the lines between science, politics, and policy are pretty fuzzy. There are many competing protocols. Are you seeing a standardization in measurement and error estimation?
- There is a need for standard methods and calculations internationally, and the IPCC plays that role globally. In the US federal system, however, the national government is letting the states be laboratories in developing these protocols, so there is less consistency in the U.S.
 - IPCC has the best available science with lots of review, so following their protocols is generally a good idea.
 - In US, parallel processes are producing some unfortunate things. For example, Chicago Climate Exchange guidelines are very lax by international standards. As another example, the CCAR has decided on a baseline methodology that is very different from UNFCCC rules and very specific to California. The baseline under CCAR is a projection of how much tree cover would be predicted based on common practice and legal and financial constraints. The UNFCCC, however, UNFCCC requires documenting how land cover has changed, using, for example, LandSat photos, then using those trends to project into the future.
 - There is a chance, under the new administration, that there will be a national program.

B. Forest regeneration

- From the work presented this week and elsewhere, it appears that to store more carbon
 in rangelands, we will need more trees, but what about limits? There are places where
 you could plant more trees but others were they've already reached their max based on
 precipitation.
- However, many places in the regions that Patrick studies in Africa, for example, are not anywhere near their maximum density. Some ethnic groups in the area do planting and regeneration activities, and in those areas you can see a doubling in the density of mature trees.

- In these areas that are seeing shifts in patterns of rainfall, such as the study areas, it may be that passive regeneration could occur (which Patrick has seen in old fields left to revert to forest) but with the replacement of the more mesic species by the drought tolerant species. So regeneration can occur, even in these areas, but the species mix will be different. It is important to note, too, that the broad trends in precipitation shifts hide more local effects, so there are actually some areas in which the rainfall has not changed much, and in those areas regeneration of mesic species may be possible.
- The extent to which regeneration is possible depends, too, on whether we want active or passive regeneration. In California, people are planting oaks, and they can grow without irrigation, but in other places, replanting forests (or planting them in areas where they did not exist previously) may require irrigation, which may not be acceptable.
- Finally, it is important to remember that while the most effective way to store carbon on rangelands is probably to plant forests, there are management activities on rangelands that could increase carbon without conversion to forest.

C. Managing for multiple uses - livestock grazing and carbon sequestration

- Groups like the California Rangeland Coalition are seeking to protect working landscapes. Can this type of science be used for groups like this to argue for protection of working landscapes?
- Working landscapes are important because eating is necessary, meaning that there is a need to manage landscapes for productive agricultural use as well as other benefits. We need to learn more about these landscapes and how to support both carbon and other productive land uses to conserve them. In both the Sahel and California, we should look at traditional agroforestry systems to learn how those systems succeed (or do not) in producing multiple benefits. We at the university focus on how the system works, but it is also necessary and worthwhile to take that information and apply it in support of goals such as conservation of working landscapes. As of now, we still do not fully understand the balance and trade-offs involved in different land uses as far as their carbon impacts.
- We've talked in the roundtable about how grazing affects carbon, but in places with
 woody plants, it seems clear that wood is better. In places where there are some woody
 species, such as oak woodlands, it seems that we could use grazing animals to help
 manage for trees for example to redistribute nutrients for tree growth. It would just be
 a question how to plan that grazing spatially and temporally to generate and credibly
 measure carbon gain for credits.
- However, adding livestock to a landscape would also mean methane emissions, so how would management practices of sequestering carbon (in soil and biomass) using livestock balance with the methane produced by the animals? When does the methane start to overwhelm the benefits?
- Livestock and methane emissions could quickly swamp out woody biomass gains.
 However, to really understand the interaction, it also matters where these livestock would have been and whether their emissions are really additional.
- We need good life-cycle assessments of these activities to better compare different land
 uses and alternatives, such as this question of livestock and trees or the issue of
 harvesting or leaving trees, as we discussed last time. The key is to keep multiple
 ecosystem services intact so that these systems are more resilient in the face of climate
 change.