Lecture 4, Characterizing the Vegetation, Part V: Phenology

- Outline
- Phenology, Defined, Quantified, Observed, Modeled
- Discuss Trends with Climate, Latitude, Global Change
- Discuss impact of phenology on agriculture, ecology, weather and climate

Phenology

- Study of the Timing of Natural Phenomena, like flowering, bud break, full leaf expansion, & leaf fall
- Derived from Greek word *phaino,* meaning to show or appear,

ESPM 129 Biometeorology 2

Phenology is a classic topic of biometeorology. The timing of many of life's events are cued by such factors as accumulated heat units and day length, things we measure and study.

Why Study Phenology?

- Determine structural and functional state of leaves, flowers and fruits
  - Know when leaves are present/absent, functioning and senescent
    - Photosynthesis, Transpiration, Albedo, VOC emissions
  - Know when flowers emerge and are pollinated
    - Fruit and reproduction

Deciduous plants operate is a quasi on/off mode, in either the deciduous and dormant stage or the leafed and photosynthesizing stage. Knowing when leaves leaf-out and how long they are active play important roles on how much carbon can be assimilated and how much water can be transpired from a vegetated landscape.

Description of principal growth stages of herbs

Stage	
0	Germination / sprouting/ bud development
1	Leaf development (main shoot)
2	Formation of side shoots / tillering
3	Stem elongation or rosette growth / shoot development (main shoot)
4	Development of harvestable vegetative plant parts (beet, tuber) / Booting (=swelling of panicle, ear, ...) (main shoot)
5	Inflorescence emergence (main shoot) / heading
6	Flowering (main shoot)
7	Development of fruit
8	Ripening of fruit and seed
9	Beginning of dormancy, die off of leaves, stem, ...

Life stages and history of an annual crop plant.

Variation in Date of Bud Burst and 75% Leaf Out by Species, at Harvard Forest



While there are geographical patterns to phenology, they can differ at a locale by species, too. Here are data from a mixed hardwood forest at Harvard Forest. Data of Richardson and O'keefe.

Hopkins Law of Phenology

- Phenology-of plants in the temperate deciduous forest biome-differs by four days for every degree of latitude, every 5 degrees of latitude and every 400 feet of altitude

Andrew Delmar Hopkins

A simple biogeographical rule, indicating how phenology, as defined by bud break or first bloom, will change with latitude and altitude.



The timing of first bloom is very distinct as one moves from south to north across the eastern half of North America. Bloom can happen as early as day 90 (end of March) in Tennessee and is as late as day 150 (end of May) in the northern tier of the US. The timing of first bloom, as determined by isolines, is very convoluted across the Rocky Mountains due to elevation.

Growing Season Length, as Detected by MODIS

S. Ganguly et al. / Remote Sensing of Environment 114 (2010) 1805–1816



Fig. 1. Mean growing season length in days for North America for the period 2001–2006 derived from the MLCD product.

One of the key phenological metrics is length of growing season. Using reflection measurements from space, scientists are able to detect the start and end of the growing season and reconstruct maps of growing season. Along the Pacific Coast and the South, growing season exceeds 300 days. Along the Mississippi River and the Corn Belt we see a short growing season. This is not the true growing season length, but the effective growing season of the annual crops planted in these regions. The Rocky Mountains and Sierra Nevada experience less than 130 days in many areas

Net Ecosystem Carbon Exchange Scales with Length of Growing Season



Baldocchi, Austral J Botany, 2008



Scientists show a strong correlation between mean annual air temperature and growing season length.



We found an interesting relationship between the start of the photosynthetic season when soil temperature cross over the mean annual air temperature. The question to be asked is if plants are in synch with their local environment. It makes some sense to start the growing season when the mean temperature is achieved, as in the spring. If leaves are launched too early there is a good chance they will experience extremely cold events. If they launch too late they miss out on important days of photosynthesis.



Using Carbon flux data we were able to id the critical day



We then applied this approach to a network of sites across the world and it works reasonably well, without tuning.

Onset of Spring is Delayed ~ 5 days with each degree reduction in mean temperature



Baldocchi et al. Int J. Biomet, 2005

To upscale this idea regionally, we plotted the zero crossing day with the mean air temperature

When Transformed onto a Climate Map, We observe a General Correspondence with N-S gradient Obtained from the denser Phenology Network



Using GIS we created a map of the spring wave.

Caveat Emptor



- Growing Season Length has More Explanatory Power across a Latitudinal Gradient than at an Individual Site
- Additional factors explaining annual NEE at a Single Site include:
  - Absence/presence winter snow
  - Occurrence of Summer Drought
  - Extent of cloudiness

Also recognize that sometimes an earlier growing season may not yield more carbon uptake. There is a growing association that years with impending drought have an early spring, but then the soil water profile is depleted sooner, which reduces photosynthesis later in the season. So an increase in annual carbon uptake does not occur.

Regional Variations in Controls on Phenology

628 W. M. JOLLY *et al.*



Sub-Tropics are Drought Deciduous

Temperate and Boreal Zones are Queued by Temperature and Daylength

One size does not fit all when describing and predicting phenology. In the warm, wet tropics, deciduousness can occur due to seasonal drought.

Drought Deciduous Forest in Thailand

http://www.ecology.kyoto-u.ac.jp/~atto/Thailand_ENG.html

Sakaerat Environmental Research Station. I was recently at the AsiaFlux meeting and was astounded to see the state of this drought deciduous tropical forest in Thailand. My prior impression was that drought was relative and only a few leaves fell.



Figure 1. (a) Definition of land surface phenological phases used in this study. (b) Simplified land cover map based on the GLC2000 Africa [*Mayaux et al.*, 2004].

Guan et al 2014 JGR Biogeosci

ESPM 129 Biometeorology

19

Timing of Drought Deciduousness across Africa. Guan et al 2014 JGR Biogeoscience



The presence and absence of leaves has a critical impact on the humidity of the air. The air is much drier without leaves. I also notice flying north south during the spring green up that on 'clear' days I don't see cumulus clouds develop over the leafless forests, but they due over those with leaves and are transpiring. A classic example of plant-atmosphere interactions and feedbacks.

Dormancy

ESPM 129 Biometeorology 21

Length of Growing season is dictated, in part by the length of the Dormant period and when that Dormant period Breaks. In this next section we will discuss methods to understand the duration of dormancy and when it breaks

Image of Steve McMillan.
http://www.sonic.net/aquatint/ART/art2Sonoma/art2pages/art2_a_fg.html

Terminology

Dormancy: is a state of reduced growth rate,
with few or no cell division in terminal or lateral meristems
(Perry, 1971 Science); Dormancy Requires a Chill Requirement

Dormancy can be Induced and Broken, Naturally or Artificially

Stages of Dormancy, Rest and Quiescence

Rest is a period when buds are dormant due to physiological conditions
Rest Ends when Chill Requirement is Fulfilled

Quiescence is a period when buds remain dormant due to unfavorable conditions; Quiescence ends when Forcing is fulfilled

Bud break and blooming are activities that are visible and signify the End of Dormancy. It is easy to measure and observe the occurrence of bud break and blooming. But what factors lead to these events and how can we quantify them? To answer these questions let's first examine Dormancy.

Processes

Temperature and Water Driven Phenology

Rest: is a function of a sum of sufficient chilling $S_{chill} = \sum_{t_1}^{t_n} R_{chill}(T)$

Quiescence: is a function of thermal forcing

Frost Hardiness: $S_{quiescence} = \sum_{t_1}^{t_n} R_{quiescence}(T)$

Bud Burst: After Cumulative Chilling and Forcing Temperatures Exceed Threshold

Bud Burst, too Early: Risk of Frost Damage

Bud Burst, too Late: Missed Opportunity for Carbon uptake

Other models are separating the Dormant period into one requiring enough chill to achieve rest, followed by enough warmth to break that rest.



At the onset of dormancy, the buds are in a resting stage. This is a period of reduced growth rate,

with few or no cell division in terminal or lateral meristems (Perry, 1971 Science). Dormancy during the Rest phase Requires a Chill Requirement, so this is the period of Chill Accumulation. Once this Chill requirement is met, the end of dormancy needs some forcing, produced by hormones and triggered by Heat requirements

I am seeing some contradiction in the literature and terminology. A number of modelers (Chuine, Kramer) speak of quiescence occurring after the Chill requirment is met. Others are calling quiescence the initial phase. TBA

Heat Units and Phenology

- Growing Degree Days, GDD
  - Sum of Average Temperature above reference
- Chill Degree Days, CDD
  - Sum of Average Temperature below Reference, e.g. 7 C and above a threshold, e.g. 0 C
- Chill Hours
- Chill Degree Hours
- Heat Degree Days

$$GDD = \sum (\frac{T_{\max} + T_{\min}}{2} - T_{ref})$$

ESPM 129 Biometeorology 25

Simple heat unit models have been popular for decades to inform nut and fruit tree breeders how much cold they need to acquire to rest and break that rest. It also instructs farmers on where and where not to plant certain cultivars and species. Growing degree days and chill degree days are the simplest. You can also use this idea for heating homes. 'days' are counted in terms of the number of degrees warmer than some reference temperature. Degrees are not counted if temperature exceeds a critical value. Average temperature is the mean between maximum and minimum temperatues.



Functional shape of simple and more complicated winter chill models. The simple chill degree method is denoted by the simple delta function. Better and more modern methods, like the Modified Utah method, penalize and subtract chill units if the temperature is too warm.



Simple models by scientists like Kramer 1994; Hanninen 1990; Chiune (1999) sum the chilling and the forcings functions based on the temperatures during the dormant seasons. Here are visual shapes of these forms.

Accumulation of Thermal Heat Units for Rest, or chill

$$R_{rest}\begin{cases} 0, & T(t) < T_{\min} \\ \dfrac{T(t)-T_{\min}}{T_{opt}-T_{\min}} & T_{\min} < T(t) < T_{opt} \\ \dfrac{T(t)-T_{\max}}{T_{opt}-T_{\max}} & T_{opt} < T(t) < T_{\max} \\ 0, & T(t) > T_{\max} \end{cases}$$



$$S_{rest} = \sum_{t_1}^{t_n} R_{rest}(T)$$

ESPM 129 Biometeorology 28

Functional Forms to define Rest.

Sources: Koen Kramer and H. Hanninen

Quiescence/Forcing



$$R_q = \begin{cases} 0, & T(t) < T_{\min} \\ \dfrac{1}{1+\exp(a_0 T(t) + b_0)}, & T(t) >= T_{\min} \end{cases}$$

ESPM 129 Biometeorology 29

Mathematical Functions and Functional shape of Forcing function

Kramer, 1994 Chuine et al 1999 PCE



Be careful with temperature sums. While they can be useful and are widely used remember, summing to sets of random numbers can yield strong coreelations, too

Trends in the Reduction in Winter Chill in CA Orchard Regions



Baldocchi and Wong, 2008 Climatic Change

ESPM 129 Biometeorology 31

Using simple Chill method, an undergraduate student, Simon Wong, and I inspected if there are trends in winter chill as California faces and is experiencing global warming. We see lots of year to year variation, but we also found trends in decreasing winter chill across the fruit growing region of California. In sufficient chill can affect flowering, flower abortion and fruit set.

The idea for this work was catalyzed by a conversation I had with my cousin, a cherry rancher, after I gave a talk at the local farm bureau. He reflected that his cherry crops were not as productive in the past because he was not getting enough winter chill. His option was to apply an expensive plant hormone to mimic the effects of chill.



Eike Luedeling, at UC Davis has spatialized past, current and future sums of winter Chill across the Central Valley of California. This is the power of GIS, to take simple point based analyzes and spatialize them and produce maps.

Winter Chill is Promoted by Valley Fog





Baldocchi and Waller, 2014 GRL

ESPM 129 Biometeorology 33

As scientists we also want to know How and Why, in addition to What. With regards to winter chill, many of us native to the Central Valley have notices we are getting less winter fog. Is this real and could it contribute to less chill. Remember with less fog, air temperatures are higher and the energy balance on the buds is greater, increasing their surface temperature. In other words, Fog keeps the buds cool, so they can accumulate chill. This is a classic example of applied biometeorology.



Why may Chill be declining? One idea is that we are experience less winter fog. Over the period of satellite remote sensing we have seen a 40% decline in the number of foggy days.

To answer this question, I collaborated with a graduate student, Eric Waller, Ph.D.. We acquired satellite data from the MODIS and AVHRR sensors. The time series goes back to 1981. We then focused on California and attempted to count the number of days each winter we could see regions of the valley in fog. We found a downward trend in the number of fog events each winter. Again lots of year to year variation with wet and dry years. But the trend is significant.

How it is Studied and Quantified

- Phenology Gardens
  - Blooming of lilac, honeysuckle or native plants
  - European Phenology Network'
    - http://www.dow.wau.nl/msa/epn/index.asp
  - National Phenology Network
    - http://www.uwm.edu/Dept/Geography/npn/
- Remote Sensing
  - Satellites
  - Digital Camera
  - Light Transmission
- Eddy Flux Measurements of Carbon Dioxide Exchange
- Models based on Thermal Time

Phenology is determined both indirectly and directly.

The longest records are observations at botantical gardens or country estates on the dates of first birds, first leaf, first flower etc. In Europe and Japan, some records span several hundred years. According to the history of phenology http://budburst.org/phenology_history the Japanese have been recording the timing of cherry blossoms for over 1200 years. Linneaus and Robert Marsham started observations in Sweden and Britain, respectively, in the 1700s.

Observations are also a great course of citizen science projects for school children.

Location of Phenology Gardens in North America



Fig. 1 Ground phenology records. Boxes show: location, center of box; number of species, box size; number of years observed at each site over the 1982–2006 study period, box color.

White et al, 2009, GCB ESPM 129 Biometeorology 37

Map of Phenology Gardens



Normalized Difference Vegetation Indices, like NDVI and EVI, enable one to study phenology from space and produce regional maps.

Light Transmission through Harvard Forest



Richardson and OKeefe ESPM 129 Biometeorology 39

Measurements of Light transmission through Forests is a good measure of dates of leaf on and leaf off.

Monitor Seasonality of Gap Fraction of Forests
with Upward Looking Camera

Gap Fraction Phenology with Upward Looking Cameras under Oak Savanna



Can Detect Start and End of Growing Season with Precision

Here is the newest and most recent data on the phenology of our oak savanna near Ione. Look how early spring was in 2016 and the range of leaf out, from the end of February to the start of April!!

With the drought leaves dropped much earlier in 2015. And the legacy effects of the drought is causing LAI to be the least this year even though we had a fairly wet spring.



Digital cameras are producing quantitative and qualitative measures of phenology. Digital information from the Red, Green Blue bands can be manipulated to produce a greeness index. Here are data from Japan by Ide et al.

In the US there is the Phenocam project, led by Andrew Richardson at Harvard. Data from cameras across the US are available at

https://phenocam.sr.unh.edu/webcam/

Our team is participating in this project

Phenology with WebCams by
Extracting Color information from
Digital Numbers of Red, Green and Blue





$$ExG = 2G - (R + B)$$

$$g_{cc} = \frac{G}{G + R + B}$$

Richardson et al., 2007, Oecologia

Digital cameras provide information on Green, Red and Blue and by assigning digital values to these pixels one can compute greenness index. By identifying a portion of the image one can take daily camera data and produce a time series.

Phenology of Flowering and Seeding Plants Complicates interpretation of Greenness Indices

Purple Flowering Alfalfa

White and Yellow flowers on Pepperweed

Rice in Seed Stage

Greenness Index over Crop, Wetland and Pasture



Saenz, Knox, Sturtevant, Koteen, Verfaillie, Baldocchi, unpublished

Cameras Monitor Management and Phenology of Crops, Rice 2013



Disced, pre-planting



Flooding, seedlings

Full canopy, vegetated

Seed Filling

Harvesting

Chopped Straw

Canopy Photosynthesis vs Vegetation Index, Rice



Phenology as a measure of Global Change

Warming in Spring, by 1 C, advances the growing season by 7 days.



Average dates of the beginning of growing season (*B*G), beginning of stem elongation for winter rye (*R*31), beginning of cherry tree blossom (*C*60) and beginning of apple tree blossom (*A*60) in Germany, 1961–2000.

Chmielewski et al (2004), AgForMet

ESPM 129 Biometeorology 49

Many long term measures of spring are showing earlier and earlier appearance

Mean annual growing season in Europe increases by 10.8 days from 1981 to 1991.



Menzel and Fabian, Nature 1999

Positive Trends in the Date of the End of the Growing Season



Dragoni and Rahman, 2012

And the end of the growing season is getting later.

Summary

- The distribution of plant functional types is associated with climatic thresholds such as minimum temperature, occurrence of frost or freezing, and water balance
- New biogeographical theories and remote sensing measurements are able to help us map the global distribution of potential and existing vegetation with greater accuracy
- Phenology concerns us with the timing of plant activities. The timing of leaf out is one of the most distinct phenological events and can cause dramatic switches in atmospheric humidity, surface energy balance and carbon uptake
- Phenology is often estimated using growing degree days
- Many new studies are showing that global warming is causing length of growing season to lengthen, hence a feedback of temperature on phenology, besides kinetics
- Timing of phenology also has a major impact of the warming and the humidification of the atmosphere and the growth of the planetary boundary layer

Estimate Degree Hours from max-min Temperatures



$$d = \tan\theta \cdot (T_{ref} - T_{\min})$$

$$\tan\theta = \frac{a}{b} = \frac{6hr}{T_{ave} - T_{\min}} = \frac{d}{c} = \frac{d}{T_{ref} - T_{\min}}$$

Chill Hours = 2 d