

[Share](#) |[Subscribe](#) [Free alerts](#) [RSS](#)[About the cover for May 2011](#)

ISSN: 0191-2917

SEARCH

Enter Keywords

- MPMI
- Phytobiomes
- Phytopathology
- Plant Disease

[search](#)[Advanced Search](#)

Resources

[Subscribe](#)[About Plant Disease](#)[First Look](#)[Most Downloaded Articles](#)[Journals Impact](#)[Submit a Manuscript](#)[Customer Care](#)[About My Password](#)[Rights and Permissions](#)[Plagiarism and Ethics](#)[Advertise](#)[e-Xtra](#)[Open Access](#)[ORCID Registry](#)

plant disease

Editor-in-Chief: Alison E. Robertson

Published by The American Phytopathological Society

[Home](#) > [Plant Disease](#) > [Table of Contents](#) > [Abstract](#)
[Previous Article](#) | [Next Article](#)May 2011, Volume 95, Number 5
Page 619<https://doi.org/10.1094/PDIS-01-11-0052>

Disease Notes

First Report of *Seiridium unicorne* Causing Bark Cankers on a Monterey Cypress in California

G. Della Rocca and **R. Danti**, CNR, Institute for Plant Protection, Area della Ricerca, Via Madonna del Piano 10, 50019 Sesto Fiorentino, Italy; and **M. Garbelotto**, Department of ESPM, 137 Mulford Hall, University of California, Berkeley 94720[Open Access](#).

In June 2009, dieback of distal branches and resin exudation associated with bark lesions were observed in an adult *Cupressus macrocarpa* tree in Sonoma County, California (Glenn Ellen; 38°21'N, 122°31'W, elevation 233 m). The fungal pathogen, *Seiridium unicorne* (Cooke and Ellis) Sutton, was obtained by plating fragments of necrotic bark from the margins of branch cankers on potato dextrose agar (PDA). Identification was based on cultural, morphological, and molecular traits (2,3). Colonies on PDA were dense, cottony, off-white at first and then turning pale gray-green, and 2.3 and 4.3 cm in diameter after 1 and 2 weeks of growth at 20°C, respectively. Colonies of the fungus showed a faster radial growth at 20°C than at 25°C. Acervuli were abundantly produced on water agar amended with autoclaved cypress seeds after 2 to 3 weeks at 18°C under a mixture of fluorescent and near UV light. Conidia were six celled (five euseptate), fusiform, 20.9 to 35.2 × 7.11 to 10.57 μm, straight or slightly curved, with four, brown median cells, and with end cells bearing unbranched appendages 2 to 5 μm long. The DNA sequence of a portion of the β-tubulin locus (GenBank Accession No. HQ678171) revealed a 100% homology with sequences of *S. unicorne* isolates from Portugal and South Africa, while being clearly distinct from sequences of *S. cupressi* and *S. cardinale* isolates (2). Greenhouse stem inoculations were performed by underbark placement of a 3-mm plug taken from the margins of a colony of the fungus grown on PDA. Inoculations were repeated twice in the spring and fall of 2010 on 10 *C. macrocarpa* saplings grown in pots for 3 years. Three months postinoculation, the pathogen could be successfully reisolated from the edges of 15 to 30 mm long elliptical lesions, present on each one of the inoculated saplings. The observed *S. unicorne* isolate is atypical because of its shorter appendages compared with the form reported in the literature (2,3). Because of its shorter conidial appendages and in vitro temperature optimum of 18 to 20°C, the fungus described here is similar to an unnamed *Coryneum* sp. observed by Wagener on *C. macrocarpa* (4). *S. unicorne* is a pathogen of many Cupressaceae in Africa, New Zealand, Japan, and some U.S. states (Georgia, South Carolina, Kansas, and Texas) (3), and although it was mentioned in a

Quick Links

[Add to favorites](#)[E-mail to a colleague](#)[Alert me when new articles cite this article](#)[Download to citation manager](#)[Related articles found in APS Journals](#)

Access provided by Univ of California

WHITE PAPER

Foundational and Translational Research Opportunities to Improve Plant Health

[Read Article Comments](#)

FOCUS ISSUE


Disease Management in the Genomics Era



Nearly 200 pages!

USDA Plant Quarantine Division report from 1963 as found on cypress in San Francisco (1), it has never been officially reported from California. Since similar disease symptoms were observed on many Cupressaceae in the course of an extensive survey performed in 2009 in California, it may be important to evaluate the relative incidence of *S. unicorn* compared with that of *S. cardinale*, a pathogen more commonly reported in association with the disease (4).

References: (1) D. F. Farr and A. Y. Rossman. Fungal Databases. Systematic Mycology and Microbiology Laboratory, ARS, USDA. Retrieved from <http://nt.ars-grin.gov/fungaldatabases/fungushost/fungushost.cfm>, 1/19/2011. (2) P. Krokene et al. Mycologia 96:1352, 2004. (3) N. A. Tisserat. Plant Dis. 75:138, 1991. (4) W. W. Wagener. J. Agric. Res. 58:1, 1939.



Cited by

AFLP analyses of California and Mediterranean populations of *Seiridium cardinale* provide insights on its origin, biology and spread pathways

G. Della Rocca, T. Osmundson, R. Danti, A. Doulis, A. Pecchioli, F. Donnarumma, E. Casalone, M. Garbelotto, and S. Woodward

Forest Pathology Jun 2013, Volume 43, Number 3, 211-221

[Crossref](#)

Impacts of exotic forest pathogens on Mediterranean ecosystems: four case studies

Matteo Garbelotto and Marco Pautasso

European Journal of Plant Pathology Dec 2011

[Crossref](#)