



NOTE

Batrachochytrium dendrobatidis: requirement for further isolate collection and archiving

Jamie Voyles^{1,*}, Katy Richards-Hrdlicka², Scott D. Cashins¹, Erica B. Rosenblum³, Alex D. Hyatt⁴, Lee Berger¹, Lee F. Skerratt¹

¹School of Public Health, Tropical Medicine and Rehabilitation Sciences, Amphibian Disease Ecology Group, James Cook University, Townsville, Queensland 4811, Australia

²School of Forestry and Environmental Sciences, Yale University, New Haven, Connecticut 06520, USA

³Department of Biological Sciences, University of Idaho, Moscow, Idaho 83844, USA

⁴Australian Animal Health Laboratory, CSIRO Livestock Industries, Geelong, Victoria 3220, Australia

ABSTRACT: The fungal pathogen *Batrachochytrium dendrobatidis* (*Bd*) causes the disease chytridiomycosis, which is lethal to many species of amphibians worldwide. Many studies have investigated the epidemiology of chytridiomycosis in amphibian populations, but few have considered possible host–pathogen coevolution. More specifically, investigations focused on the evolution of *Bd*, and the link with *Bd* virulence, are needed. Such studies, which may be important for conservation management of amphibians, depend on access to *Bd* isolates. Here we provide a summary of known *Bd* isolates that have been collected and archived in various locations around the world. Of 257 *Bd* isolates, we found that 53 % originate from ranids in the United States. In many cases, detailed information on isolate origin is unavailable, and it is unknown how many isolates are cryo-archived. We suggest the creation of a centralized database of isolate information, and we urge researchers and managers to isolate and archive *Bd* to facilitate future research on chytridiomycosis.

KEY WORDS: Amphibian declines · *Batrachochytrium dendrobatidis* · Chytridiomycosis · Pathogen preservation · Wildlife disease

Resale or republication not permitted without written consent of the publisher

INTRODUCTION

Chytridiomycosis is caused by the fungal pathogen *Batrachochytrium dendrobatidis* (*Bd*; Berger et al. 1998, Longcore et al. 1999). In infected amphibians, *Bd* is found in the superficial layers of the epidermis and disrupts normal osmoregulatory functioning in the skin (Berger et al. 1998, Voyles et al. 2007, 2009a). Mass mortality events have coincided with the appearance of *Bd* in wild amphibian populations (Berger et al. 1998, Lips et al. 2006). The effect on some host species is extreme, leading to dramatic declines and possibly extinctions (Schloegel et al. 2006, Skerratt et al. 2007). Some populations that survive initial declines persist with various levels of infection (Retallick et al. 2004, Woodhams & Alford 2005), and *Bd* maintains at least

moderate virulence in some species many years after introduction (Murray et al. 2009). These observations may be explained by factors such as variability in host resistance (Woodhams et al. 2007), host behavior (Rowley & Alford 2007), or environmental characteristics (James et al. 2009), but shifts in *Bd* virulence are also plausible.

Some evidence supports the possibility of differential virulence among *Bd* isolates. Laboratory experiments suggest that *Bd* virulence differs among isolates when introduced to a single susceptible amphibian species (Berger et al. 2005, Retallick & Miera 2007). In addition, phenotypic differences among isolates in proteomic signatures, morphological characteristics, and zoospore production (Fisher et al. 2009) could be associated with differences in virulence, although the mol-

*Email: jamie.voyles@gmail.com

ecular basis for any differences in isolate virulence has not yet been identified. There seems to be relatively low genetic variability among isolates collected from globally widespread sources (Morehouse et al. 2003, James et al. 2009), but a recent study reported differential virulence among *Bd* strains that were either endemic to Japanese native amphibians or associated with introduced species (Goka et al. 2009). The possibility of differential virulence among distinct isolates highlights the requirement for ongoing surveillance, continued development of diagnostic assays for *Bd*, and further virulence research. However, advances in chytridiomycosis research will require access to *Bd* isolates. The methods for isolating and purifying *Bd* were first established in 1999 (Longcore et al. 1999). Some *Bd* isolates have been cryo-archived for future research (Boyle et al. 2003), and many are actively passaged under different nutrient and temperature conditions. Here we review available information on known *Bd* isolates.

MATERIALS AND METHODS

Isolate records were gathered from personal lists (L. Berger, James Cook University; A. Hyatt, CSIRO Australian Animal Health Laboratories; J. Longcore, University of Maine) and peer-reviewed papers (Morehouse et al. 2003, Berger et al. 2005, Morgan et al. 2007, Rosenblum et al. 2008, Symonds et al. 2008, Fisher et al. 2009, James et al. 2009). The database is comprised of information for 257 isolates, which probably represent a subset of all existing isolates (i.e. information on additional isolates may not be currently available in the published literature). We collected as much information as possible on isolate origin (host species, life stage, location, and disease status), isolate storage history (passage history, location, current storage conditions), and contact information for researchers working with isolates.

RESULTS

The majority of *Bd* isolates, approximately 53%, originate from ranids in the USA. However, a disproportionate number of originate from *Rana muscosa* (44 isolates) and *R. sierrae* (57 isolates) that were collected for a population genetics study in California (Morgan et al. 2007). Only 2 isolates come from caudates (Fig. 1). Most of the isolates (156) are from amphibian populations in the United States, which again is a reflection of the large number of isolates collected from *R. muscosa* and *R. sierrae* in California (Morgan et al. 2007).




Fig. 1. *Batrachochytrium dendrobatidis*. Number of isolates per amphibian family collected and maintained in laboratory collections

Additional information on these isolates such as isolate origin (host species, date, life stage, location, disease status) and isolate storage history (passage history, location, current storage conditions) was collected whenever possible. Detailed information was difficult to obtain from the published literature, and in some cases no details were available. Of the identified isolates, little to no information was available about which isolates have been cryo-archived and are thus available for future research. Additionally, our analysis of available *Bd* isolates suggests that very few isolates are being collected and archived from important regions where *Bd*-associated amphibian declines have occurred, or may be currently taking place (e.g. Central America: Republic of Panama, Woodhams et al. 2008; Montserrat, G. Garcia pers. comm.; Southeast Asia: Indonesia, Kusriani et al. 2008; Philippines, R. Brown pers. comm.).

DISCUSSION

Confronting disease-related declines requires addressing a specific set of problems: identifying the etiological agent and possible point of origin, developing diagnostic assays and sampling protocols, and understanding the mechanisms of pathogenesis, transmission, and evolution of host–pathogen dynamics. Many of these challenges have been successfully addressed (e.g. Berger et al. 1998, Boyle et al. 2003, Hyatt et al. 2007, James et al. 2009, Voyles et al. 2009a), but others are still being investigated. To facilitate projects that will meet these challenges, we recommend that additional *Bd* isolates be collected and cryo-archived.

Table 1. Summary of *Batrachochytrium dendrobatidis* (*Bd*) isolates. Isolate records were collected from published literature and personal lists. Host species refers to the amphibian species from which *Bd* was originally isolated

Host	Total	Host	Total
Canada	12	Brazil	1
<i>Rana catesbeiana</i>	12	<i>Rana catesbeiana</i>	1
USA	156	Peru	1
<i>Ambystoma tigrinum</i>	1	Unknown species	1
<i>Bufo americanus</i>	1	Venezuela	2
<i>Bufo boreas</i>	4	<i>Rana catesbeiana</i>	2
<i>Bufo haematiticus</i>	1	South America (country unknown)	3
<i>Dendrobates auratus</i>	1	<i>Cochranella albomaculata</i>	1
<i>Dendrobates azureus</i>	5	<i>Colostethus flotator</i>	1
<i>Dyscophus guineti</i>	1	<i>Hyalinobatrachium colymbiphylum</i>	1
<i>Hyla arenicolor</i>	1	Puerto Rico	1
<i>Litoria caerulea</i>	1	<i>Eleutherodactylus coqui</i>	1
<i>Rana aurora draytonii</i>	1	Ghana	3
<i>Rana catesbeiana</i>	7	<i>Xenopus tropicalis</i>	3
<i>Rana clamitans</i>	4	South Africa	7
<i>Rana muscosa</i>	44	<i>Afrana fuscigula</i>	3
<i>Rana palustris</i>	1	<i>Xenopus laevis</i>	4
<i>Rana pipiens</i>	9	Africa (country unknown)	12
<i>Rana sierrae</i>	57	<i>Xenopus laevis</i>	12
<i>Rana sylvatica</i>	1	Japan	1
<i>Rana yavapaiensis</i>	12	<i>Ceratophrys cranwelli</i>	1
<i>Xenopus tropicalis</i>	1	Australia	15
Unknown species	3	<i>Limnodynastes dumerilii</i>	1
Panama	25	<i>Limnodynastes peronii</i>	1
<i>Atelopus zeteki</i>	2	<i>Litoria booroolongensis</i>	1
<i>Bufo haematiticus</i>	1	<i>Litoria caerulea</i>	2
<i>Centrolenella ilex</i>	1	<i>Litoria lesueuri</i>	1
<i>Cochranella euknemos</i>	1	<i>Litoria peronii</i>	1
<i>Colostethus inguinalis</i>	3	<i>Litoria rheocola</i>	2
<i>Colostethus nubicola</i>	1	<i>Mixophyes fasciolatus</i>	3
<i>Eleutherodactylus bufoniformis</i>	1	<i>Mixophyes fleayi</i>	2
<i>Eleutherodactylus caryophyllaceus</i>	2	<i>Nyctimystes dayi</i>	1
<i>Eleutherodactylus museosus</i>	2	Spain	9
<i>Eleutherodactylus podi-noblei</i>	1	<i>Alytes muletensis</i>	6
<i>Eleutherodactylus talamancae</i>	2	<i>Alytes obstetricans</i>	3
<i>Hyla pameri</i>	1	UK	1
<i>Phyllomedusa lemur</i>	1	<i>Triturus vulgaris</i>	1
<i>Smilisca phaeota</i>	4	Unknown country	8
<i>Smilisca sila</i>	2	Unknown species	8
		GRAND TOTAL	257

Priority targets for *Bd* collection and cryo-preservation have been suggested (Voyles et al. 2009b), and recording detailed information on any additional isolates will be important for basic disease research.

Although protocols for isolating (Longcore et al. 1999) and cryo-archiving (Boyle et al. 2003) *Bd* are readily available, some of the required skills are not standard for medical microbiology labs. Obtaining isolates can be difficult, requiring some technical skills and persistence. A tutorial on basic techniques is available in multiple languages (see www.bdbank.org) and can be used in conjunction with the published literature, regional training courses, and consultation with a World Organization for Animal Health (OIE) reference laboratory for chytridiomycosis (e.g. the Australian Animal Health Laboratory, Geelong). The Bdbank

website provides an online forum to facilitate virulence research on *Bd*. To that end, we hope to collate and standardize data on global isolates into a single database where information can be shared and accessed by chytridiomycosis researchers. Although isolate information can be cataloged on this website, there is, as yet, no formal arrangement for physical storage of isolates. Isolates can be archived at any laboratory, but for long-term storage and access, isolates should also be cryo-archived at the Australian Animal Health Laboratory, Geelong, Victoria.

Acknowledgements. We thank J. E. Longcore and D. Boyle for their contributions on isolate information. We also thank R. Brown, G. Garcia, J. Lopez, and Q. Bloxam for providing information on frog declines.

LITERATURE CITED

- Berger L, Speare R, Daszak P, Green DE and others (1998) Chytridiomycosis causes amphibian mortality associated with population declines in the rain forests of Australia and Central America. *Proc Natl Acad Sci USA* 95: 9031–9036
- Berger L, Marantelli G, Skerratt LF, Speare R (2005) Virulence of the amphibian chytrid fungus *Batrachochytrium dendrobatidis* varies with the strain. *Dis Aquat Org* 68: 47–50
- Boyle DG, Hyatt AD, Daszak P, Berger L and others (2003) Cryo-archiving of *Batrachochytrium dendrobatidis* and other chytridiomycetes. *Dis Aquat Org* 56:59–64
- Fisher MC, Bosch J, Yin Z, Stead DA and others (2009) Proteomic and phenotypic profiling of the amphibian pathogen *Batrachochytrium dendrobatidis* shows that genotype is linked to virulence. *Mol Ecol* 18:415–429
- Goka K, Yokoyama J, Une Y, Kuroki T and others (2009) Amphibian chytridiomycosis in Japan: distribution, haplotypes, and possible route of entry into Japan. *Mol Ecol* 18: 4757–4774
- Hyatt AD, Boyle DG, Olsen V, Boyle DB and others (2007) Diagnostic assays and sampling protocols for the detection of *Batrachochytrium dendrobatidis*. *Dis Aquat Org* 73:175–192
- James TY, Litvintseva AP, Vilgalys R, Morgan JAT and others (2009) Rapid global expansion of the fungal disease chytridiomycosis into declining and health amphibian populations. *PLoS Pathog* 5:e1000458, doi:10.1371/journal.ppat.1000458
- Kusriani MD, Skerratt LF, Garland S, Berger L, Enderwin W (2008) Chytridiomycosis in frogs of Mount Gede Pangrango, Indonesia. *Dis Aquat Org* 82:187–194
- Lips KR, Brem F, Brenes R, Reeve JD and others (2006) Emerging infectious disease and the loss of biodiversity in a Neotropical amphibian community. *Proc Natl Acad Sci USA* 103:3165–3170
- Longcore JE, Pessier AP, Nichols DK (1999) *Batrachochytrium dendrobatidis* gen. et sp. nov., a chytrid pathogenic to amphibians. *Mycologia* 91:219–227
- Morehouse EA, James TY, Ganley ARD, Vilgalys R, Berger L, Murphy PJ, Longcore JE (2003) Multilocus sequence typing suggests the chytrid pathogen of amphibians is a recently emerged clone. *Mol Ecol* 12:395–403
- Morgan JAT, Vredenburg VT, Rachowicz LJ, Knapp RA and others (2007) Population genetics of the frog-killing fungus *Batrachochytrium dendrobatidis*. *Proc Natl Acad Sci USA* 104:13845–13850
- Murray KA, Skerratt LF, Speare R, McCallum H (2009) Impact and dynamics of disease in species threatened by the amphibian chytrid fungus, *Batrachochytrium dendrobatidis*. *Conserv Biol* 23:1242–1252
- Retallick RWR, Miera V (2007) Strain differences in the amphibian chytrid *Batrachochytrium dendrobatidis* and non-permanent, sub-lethal effects of infection. *Dis Aquat Org* 75:201–207
- Retallick RWR, McCallum H, Speare R (2004) Endemic infection of the amphibian chytrid fungus in a frog community post-decline. *PLoS Biol* 2:e351, doi:10.1371/journal.pbio.0020351
- Rosenblum EB, Stajich JE, Maddox N, Eisen MB (2008) Global gene expression profiles for life stages of the deadly amphibian pathogen *Batrachochytrium dendrobatidis*. *Proc Natl Acad Sci USA* 105:17034–17039
- Rowley JJJ, Alford RA (2007) Behaviour of Australian rainforest stream frogs may affect the transmission of chytridiomycosis. *Dis Aquat Org* 77:1–9
- Schloegel LM, Hero JM, Berger L, Speare R, McDonald K, Daszak P (2006) The decline of the sharp-snouted day frog (*Taudactylus acutirostris*): the first documented case of extinction by infection in a free-ranging wildlife species? *EcoHealth* 3:35–40
- Skerratt LF, Berger L, Speare R, Cashins S and others (2007) Spread of chytridiomycosis has caused the rapid global decline and extinction of frogs. *EcoHealth* 4:125–134
- Symonds EP, Trott DJ, Bird PS, Mills P (2008) Growth characteristics and enzyme activity in *Batrachochytrium dendrobatidis* isolates. *Mycopathologia* 166:143–147
- Voyles J, Berger L, Young S, Speare R and others (2007) Electrolyte depletion and osmotic imbalance in amphibians with chytridiomycosis. *Dis Aquat Org* 77:113–118
- Voyles J, Young S, Berger L, Campbell C and others (2009a) Pathogenesis of chytridiomycosis, a cause of catastrophic amphibian declines. *Science* 326:582–585
- Voyles J, Cashins SD, Rosenblum EB, Puschendorf R (2009b) Preserving pathogens for wildlife conservation: a case for action on amphibian declines. *Oryx* 43:527–529
- Woodhams DC, Alford RA (2005) Ecology of chytridiomycosis in rainforest stream frog assemblages of tropical Queensland. *Conserv Biol* 19:1449–1459
- Woodhams DC, Rollins-Smith LA, Alford RA, Simon MA, Harris RN (2007) Innate immune defenses of amphibian skin: antimicrobial peptides and more. *Anim Conserv* 10: 425–428
- Woodhams DC, Kilburn VL, Reinert LK, Voyles J and others (2008) Chytridiomycosis and amphibian population declines continue to spread eastward in Panama. *EcoHealth* 5:268–274

Editorial responsibility: Cynthia Carey,
Boulder, Colorado, USA

Submitted: September 2, 2009; Accepted: February 10, 2010
Proofs received from author(s): April 23, 2010