



ARTICLE

Mega-disturbances cause rapid decline of mature conifer forest habitat in California

Zachary L. Steel¹  | Gavin M. Jones^{2,3}  | Brandon M. Collins^{1,4} |
 Rebecca Green⁵ | Alexander Koltunov^{6,7} | Kathryn L. Purcell⁸ |
 Sarah C. Sawyer⁹ | Michèle R. Slaton⁶ | Scott L. Stephens¹ | Peter Stine¹⁰ |
 Craig Thompson¹¹

¹University of California, Berkeley, California, USA

²USFS Rocky Mountain Research Station, Albuquerque, New Mexico, USA

³University of New Mexico, Albuquerque, New Mexico, USA

⁴USFS Pacific Southwest Research Station, Davis, California, USA

⁵Sequoia & Kings Canyon National Park, Three Rivers, California, USA

⁶USFS Pacific Southwest Region, McClellan, California, USA

⁷University of California, Davis, California, USA

⁸USFS Pacific Southwest Research Station, Coarsegold, California, USA

⁹USFS Pacific Southwest Region, Vallejo, California, USA

¹⁰Stine Wildland Resources Science, Sacramento, California, USA

¹¹USFS Pacific Southwest Research Station, Fresno, California, USA

Correspondence

Zachary L. Steel

Email: zsteel@berkeley.edu

Funding information

California Department of Forestry and Fire Protection, Grant/Award Number: #8GG21823

Handling Editor: Carolyn Hull Sieg

Abstract

Mature forests provide important wildlife habitat and support critical ecosystem functions globally. Within the dry conifer forests of the western United States, past management and fire exclusion have contributed to forest conditions that are susceptible to increasingly severe wildfire and drought. We evaluated declines in conifer forest cover in the southern Sierra Nevada of California during a decade of record disturbance by using spatially comprehensive forest structure estimates, wildfire perimeter data, and the eDaRT forest disturbance tracking algorithm. Primarily due to the combination of wildfires, drought, and drought-associated beetle epidemics, 30% of the region's conifer forest extent transitioned to nonforest vegetation during 2011–2020. In total, 50% of mature forest habitat and 85% of high density mature forests either transitioned to lower density forest or nonforest vegetation types. California spotted owl protected activity centers (PAC) experienced greater canopy cover decline (49% of 2011 cover) than non-PAC areas (42% decline). Areas with high initial canopy cover and without tall trees were most vulnerable to canopy cover declines, likely explaining the disproportionate declines of mature forest habitat and within PACs. Drought and beetle attack caused greater cumulative declines than areas where drought and wildfire mortality overlapped, and both types of natural disturbance far outpaced declines attributable to mechanical activities. Drought mortality that disproportionately affects large conifers is particularly problematic to mature forest specialist species reliant on large trees. However, patches of degraded forests within wildfire perimeters were larger with greater core area than those outside burned areas, and remnant forest habitats were more fragmented within burned perimeters than those affected by drought and beetle mortality alone. The percentage of mature forest that survived and potentially benefited from lower severity wildfire increased over time as the total extent of mature forest declined. These areas provide some opportunity for improved resilience to future disturbances, but strategic management interventions are likely also necessary to

mitigate worsening mega-disturbances. Remaining dry mature forest habitat in California may be susceptible to complete loss in the coming decades without a rapid transition from a conservation paradigm that attempts to maintain static conditions to one that manages for sustainable disturbance dynamics.

KEYWORDS

climate change, drought, fisher, forest conservation, forest disturbance, habitat loss, spotted owl, wildfire

INTRODUCTION

Mature forests characterized by large, old trees are essential ecosystems that support biodiversity conservation and ecological function globally (Lindenmayer & Laurance, 2017). Additionally, these forests provide high levels of terrestrial carbon storage (Stephenson et al., 2014) that when managed in accordance with their natural disturbance regime, can be strong carbon sinks (Liang et al., 2018). Mature forests in the western USA, particularly those composed of tall, old trees and complex understories, support numerous species of conservation concern including the spotted owl (*Strix occidentalis*) and fisher (*Pekania pennanti*) (North et al., 2017; Purcell et al., 2009). Conserving the habitat for these species while also managing to reduce wildfire hazard has been a major objective of dry forest management for decades (Stephens et al., 2019; Thompson et al., 2011). Historically, mature forest patches characterized by large fire-resistant trees and varying densities made up a substantial component of the landscape vegetation mosaic (Hessburg et al., 2015; Lydersen & Collins, 2018) and were maintained by frequent, low-to-moderate severity fire ignited by lightning and Indigenous peoples (Anderson, 2013; van Wagtenonk et al., 2018).

Through a combination of past logging and increasingly severe ecological disturbances, mature forests around the world are in decline (Lindenmayer et al., 2012). In the western USA, past timber harvesting that focused on large-tree removal is primarily responsible for the limited contemporary extent of mature forest habitat (Collins et al., 2017). Further, widespread fire suppression and exclusion of Indigenous burning during the 20th and 21st centuries has allowed small fire-sensitive trees and shrubs to replace the previously extracted large fire-resistant trees (Bernal et al., 2022; Knapp et al., 2013; Taylor, 2004). These changes to forest structure and flammability plus a warming climate have led to increasingly severe disturbance cycles that may pose an existential threat to spatially limited, remnant mature forests (Figure 1; Steel et al., 2015; Stephens et al., 2018). Patch size of forest loss due to severe wildfire is also increasing, whereas unburned refugia within fire perimeters are becoming more

fragmented and sustained in smaller patches (Steel et al., 2018), which can lead to slow or unsuccessful re-establishment of conifer species dependent on live seed trees (Welch et al., 2016). Forest densification can also lead to extensive tree mortality during drought due to a combination of elevated water stress and associated beetle attack (Fettig et al., 2019). This is particularly impactful for mature forests when beetle species preferentially select large host trees, such as sugar pine (*Pinus lambertiana*) and ponderosa pine (*P. ponderosa*) in the Sierra Nevada (Stephenson et al., 2019).

These recent and increasingly extensive losses of mature forests threaten the persistence of wildlife species that require a mosaic of seral stages, including stands of large trees and multilayered canopies. Interiors of large high severity burn patches constitute poor habitat for mature forest specialists and support lower biodiversity for at least some taxa (Jones et al., 2020; Steel, Fogg, et al., 2022). Even black-backed woodpeckers, long held up as an example of fire-dependent species in dry western forests (Hutto, 2008; Odion & Hanson, 2013), have recently been shown to be sensitive to fire patch size with large, homogenous high severity patches creating sink habitats for juveniles (Stillman et al., 2019, 2021). Further, the loss of fire refugia limits the capacity of many species to recolonize the postfire landscape (Cunningham et al., 2003; Thompson et al., 2021; Vanbianchi et al., 2017).

Several factors have led to documented or likely declines in populations of mature forest-dependent species in the western USA, including the California spotted owl and the West Coast population of fishers, with projected further loss of habitat posing a grave threat to these populations (Aubry et al., 2013; Gutiérrez et al., 2017; USFWS, 2020). The southern Sierra Nevada in particular supports populations at risk of climate-related shifts in drought and wildfire patterns, including the federally endangered Southern Sierra Nevada Distinct Population Segment of fishers. The region's complex topography and elevation range also provides potential climate refugia for these species, if such habitats avoid catastrophic disturbance. Policies adopted to protect remnant patches of mature forest habitat have been in place for several




FIGURE 1 Conceptual figure showing trends in areas of conifer forests and of mature forest habitat under regimes of (a) historic, (b) current, and (c) potential future dynamics (achieved via “managed dynamics”). The late 2010s and early 2020s have been characterized by compounding mega-disturbances and extensive declines of mature forest habitat. Future managed dynamics include the restoration of small, frequent, low intensity disturbances that can help stabilize and perhaps reverse these recent trends. Vertical gray bars represent hypothetical disturbance events, with thicker bars indicating greater size/severity.

decades (Gutiérrez et al., 2017). Implementing such policies has led to land management designations such as spotted owl protected activity centers (PACs) in the Sierra Nevada National Forest Plans, and late-successional reserves in the Northwest Forest Plan. Spotted owl PACs are a roughly 120-ha land allocation designated wherever territorial owls are found in the Sierra Nevada portion of the California spotted owl range and delineated to include the best habitat (i.e., multistory, large-tree, and dense canopy cover habitat). Late-successional reserves are land allocations, representing 30% of the federal land within the range of northern spotted owls, designed to serve as habitat for late-successional- and old-growth-related species. Although these policies have effectively ended the harvest of large, old trees on United States Forest Service lands, they also limit other management activities (e.g., thinning from below, prescribed fire) making restoration of these disturbance-adapted forests more difficult (Collins et al., 2010). Furthermore, these policies implicitly assume that “protected” habitat areas can be preserved, more or less in their current form, in perpetuity (USDA, 2019). Recent disturbance trends in western forests create a test of this assumption and of the efficacy of a static approach to habitat conservation in disturbance-prone systems. Results from the Pacific Northwest suggest that in dynamic, disturbance-dependent forests, this assumption is not well supported (Davis et al., In Press). Under climate change, a static approach to mature forest conservation may be even less effective in drier and warmer regions such as the southern Sierra Nevada.

In this paper, we describe the change in the extent of mature conifer forests in the southern Sierra Nevada of

California during 2011–2020, a decade and ecoregion characterized by compounding severe wildfires and drought follow prolonged fire exclusion. The primary objectives of this analysis are to: (1) quantify the decline of conifer forest cover and of mature forest habitat due to drought (and associated beetle attack) and wildfire between 2011 and 2020; (2) test the assumption that static areas of restricted management will conserve mature forest habitat over time by examining whether rates of decline differed within versus outside of spotted owl protected areas; (3) compare the spatial pattern of drought/beetle-caused mortality with wildfire effects to better understand their differential impacts on mature forest habitat and the species that depend on it; and (4) assess whether and where forests have survived recent wildfires, potentially improving their resilience to future severe disturbance. Finally, considering the overall magnitude and extent of these disturbances, we discuss a potential shift in approach toward managing habitat for sensitive species, from one dominated by mega-disturbances toward a restoration of smaller, frequent disturbances necessary for the persistence of mature forests in fire- and drought-adapted ecosystems (Figure 1).

METHODS

Study area

Our study area is defined by the southern Sierra Nevada ranging approximately from 35° to 39° latitude and –121° to –117° longitude (Figure 2). We focus on forests




FIGURE 2 Map of forest and mature forest habitat transitions due to drought and wildfire mortality between 2011 and 2020. (a) Areas of conifer forest persistence and transitions in the southern Sierra Nevada. (b) An example of the spatial pattern of transition where drought and beetle infestation led to relatively small, but widely dispersed, patches of mortality. (c) An example of where the combination of drought and wildfire led to larger and more aggregated patches of mortality (i.e., within the 2020 Castle fire). Forests and mature forest (moderate and high density) habitats were considered to have experienced transition if canopy cover had declined below 25% and 40%, respectively. Areas initially considered mature were further limited to forests where the average height of the tallest trees exceeded 30 m. Black polygons show the perimeters of medium to large (≥ 5000 ha) wildfires during the study period and areas in the Sierra Nevada not dominated by conifer species are shown in gray.

that were conifer dominated as of 2011. Hardwood-dominated forests are also an important habitat for mature forest specialists (Aubry et al., 2013; Green et al., 2019; North et al., 2000). However, we limit our analysis to conifer-dominated forests because the tree mortality data used (described below) have not yet been

calibrated for nonconifer cover types. At lower elevations, conifer forests are dominated by ponderosa pine (*Pinus ponderosa*), white fir (*Abies concolor*), and incense cedar (*Calocedrus decurrens*), transitioning to a mixed-conifer community at mid elevations that also includes sugar pine (*Pinus lambertiana*), Douglas-fir

(*Pseudotsuga menziesii*), giant sequoia (*Sequoiadendron giganteum*), and California black oak (*Quercus kelloggii*) (Mayer & Laudenslayer, 1988; North et al., 2016). At higher elevations, Jeffrey pine (*Pinus jeffreyii*), red fir (*A. magnifica*), lodgepole pine (*Pinus contorta* var. *murrayana*), and western white pine (*Pinus monticola*) become dominant (Mayer & Laudenslayer, 1988; North et al., 2016). The southern Sierra Nevada has a Mediterranean-type climate where most precipitation occurs between November and May followed by an annual dry period broken up by sporadic summer thunderstorms (North et al., 2016).

Prior to Euro-American colonization, fire return intervals were short, especially in ponderosa pine and mixed-conifer forests (Safford & Stevens, 2017). Fire severity in these forests was predominantly low to moderate, resulting in the survival of most mature fire-resistant trees (Safford & Stevens, 2017). Return intervals were longer and the share of high severity effects (i.e., >95% vegetation mortality; Miller et al., 2009) were somewhat higher in higher elevation red fir and subalpine forest (Meyer & North, 2019). Due to 20th and 21st centuries fire suppression and exclusion, contemporary return intervals are much longer than their historic reference, although fire frequency and severity have both increased markedly in recent decades due to fuel accumulation, human ignitions, and climate change (Steel et al., 2015; Westerling, 2016; Williams et al., in revision). Severe drought was not uncommon in this region during the last two millennia (Swetnam, 1993). However, the most recent drought (2012–2016) may have been the most severe event in the last 1200+ years (Robeson, 2015).

Data and analysis

We used F³ data of canopy cover and large-tree height to map conifer forest and mature conifer forest cover as of 2011 (Huang et al., 2018). The F³ product integrates Forest Inventory and Analysis (FIA) field data with landscape and vegetation succession models to generate spatially contiguous estimates of stand structure, which has been validated for Sierra Nevada forests (Huang et al., 2018). We defined conifer forests as areas with a minimum of 25% canopy cover dominated by conifer species. This threshold is consistent with the USDA’s vegetation classification system in which tree-dominated vegetation with a minimum height of 5 m and 25%–100% canopy closure is considered forest or woodland (Brewer et al., 2005). Here, 10%–25% tree canopy cover can be considered sparse woodland or savanna vegetation. We defined mature forests as a subset of conifer forests in which the average height of the 40 largest diameter trees

within the area equivalent to a FIA plot (0.4 ha) is at least 30 m. Cover of tall trees is particularly valuable for mature forest specialists such as the spotted owl (North et al., 2017). We further classified mature forests into subgroups using the F³ data and following the California Wildlife Habitat Relationships classification of moderate density (40%–59% canopy cover) and high density (≥60% canopy cover) forests, which support various life history requisites (foraging, nesting/denning) and demographic parameters (reproduction, survival) of mature forest wildlife (Table 1). For example, 56%–61% cover may be optimal for fisher resting sites (Purcell et al., 2009) and California spotted owls tend to occupy nest sites with >70% canopy cover (Tempel et al., 2016). Forests characterized by tall trees, but <40% canopy cover may be considered low density mature forests. This forest type only covered 1076 ha in 2011, which constituted 1% of mature forest extent and 0.1% of overall conifer forest extent. The savanna and low density mature forest subclasses were not analyzed independently here as our focus was on species associated with higher canopy covers.

Estimates of live canopy cover decline (at 30 m resolution) were obtained from the Ecosystem Disturbance and Recovery Tracker (eDaRT), an image analysis system that processes all usable historic Landsat imagery (normally at the 8- or 16-day step). The eDaRT algorithm statistically models normal (e.g., phenology driven) variability in multiple vegetation indices across the image time series, and tracks model residuals through time to detect disturbance events as anomalous changes in the residuals’ trajectories, relative to a recent baseline (Koltunov et al., 2015, 2020). For detected disturbances, the algorithm estimates the resulting loss of live tree canopy cover (as pixel area fraction) using the eDaRT Mortality Magnitude Index (MMI). In this paper, we used the MMI values integrated on an annual basis between 2011 and 2020 (detailed description

TABLE 1 Conifer forest classification definitions.

Classification	Canopy cover range (%)	Large-tree height minimum (m)
Nonforest	0–25	None
Sparse woodland/savanna ^a	10–25	5
Conifer forest	25–100	5
Mature forest	40–100	30
Low density ^a	25–40	30
Moderate density	40–60	30
High density	60–100	30

Note: Classifications are hierarchical with mature forests being a subset of conifer forest and mature forests further subdivided into density classes.

^aThe savanna and low density mature subclasses are not explicitly analyzed.

in Appendix S1). Conifer forests and mature forests are considered to have transitioned to a different vegetation class if live canopy cover drops below their minimum canopy cover thresholds. Although increases in live canopy cover did occur in some instances over this time period, this new cover was predominated by seedling to sapling size trees following large-tree mortality events (Young et al., 2020). Thus, only declines in canopy cover were considered here.

The current eDaRT version does not directly differentiate the cause of detected disturbances (e.g., fire vs. drought and beetle infestation), thus necessitating additional analyses to attribute detected disturbance to tree mortality. Specifically, we assumed eDaRT-detected reduction in canopy within wildfire perimeters of a given year to be attributable to a combination of wildfire and underlying drought conditions, and within management areas to be a combination of timber harvest or fuels reduction treatments and drought conditions. To delineate burned areas for each study year, we used California's interagency fire perimeter database (available at <https://frap.fire.ca.gov/frap-projects/fire-perimeters/>; FRAP, 2022). To delineate areas managed for timber harvest or fuels reduction treatments we used the Knight et al. (2022) dataset that combined spatially explicit activities data from the California Department of Forestry and Fire Protection (CalFire) and the United States Forest Service (FACTS database); and crosswalked these activities by treatment type and intensity. Knight et al. (2022) also refined the spatial and temporal accuracy of the combined data using satellite imagery and the Continuous Change Detection and Classification algorithm. While these data include a wide range of activity types, we limited our assessment to activity types that can significantly reduce live canopy cover at the Landsat pixel scale (e.g., clear-cuts and thinning). This dataset does not include activities on National Parks land, where mechanical tree removal is minimal and at the time of writing does not include 2020 activities. Canopy declines in areas and years in which no fire or management activity occurred were attributed to drought and associated beetle kill only. This assumption likely resulted in a slight overestimate of drought mortality in 2020 when management activity data were not available.

Spotted owl PAC polygons were obtained from a United States Forest Service geospatial dataset and included a total of 651 PACs in our study area, 96% of which were established before 2011. A regular grid of non-PAC sample points was generated across our study area but limited to Forest Service lands and outside established PACs. Grid spacing (3.9 km) was set to allow an equal number of non-PAC points (651) within the 2011 extent of conifer forests. Sample points were

buffered to create non-PAC areas equal to the mean size of study area PACs (128 ha). The 2011 canopy cover, 2011 average large-tree height, and relative decline in canopy cover between 2011 and 2020 were summarized for each PAC, all non-PAC Forest Service lands, and for each non-PAC sample area. Relative canopy cover declines were calculated as the percentage of 2011 canopy cover that was lost between 2011 and 2020. Mean values within each PAC and non-PAC area were used for statistical comparison. To assess whether forest habitats within PACs were successfully preserved despite regional disturbances we modeled relative decline in cover as a function of sample class (PAC or non-PAC). Separately, we modeled relative decline in cover as a function of 2011 canopy cover and 2011 average large-tree height. Structural and PAC class variables were not combined in the same model to avoid statistical confounding as PAC delineation is determined largely by forest structural characteristics (USDA, 2004). We fitted generalized linear models using a beta error structure with a logit-link because relative canopy cover decline is a proportion bounded between zero and one. Models were fitted using the *BRMs* package in program R (Bürkner, 2017; R Core Team, 2021).

We compared landscape pattern within and outside burned areas to contrast the effect of the combined impacts of wildfire and drought/beetles with areas only affected by drought and beetle attack. Patches of all forest and mature forest that transitioned to lower density classes between 2011 and 2020 were assessed for size and core area. Area ≥ 90 m (3 pixels) from nontransitioned edge was defined as core area, which approximated distance estimates of when conifer seed dispersal and habitat use of some forest-associated species become minimal (i.e., 100 m; Jones et al., 2020; Kramer et al., 2021). Additionally, to assess the degree of fragmentation due to each combination of disturbances, we calculated the aggregation index of forest cover prior to the disturbances in question (2011), as well as within and outside of burned areas at the end of our study period in 2020. Assessments of landscape pattern were conducted for each forest class. We focused this analysis on drought and wildfire patterns only because the extent of mechanical management activities during our study period was relatively minimal (see “*Results*”).

To assess the degree to which forests survived a recent fire and potentially experienced partial restoration of the historic fire regime, we calculated years since the last fire for all areas remaining as forest and mature forest each year. Forested pixels were considered to have experienced recent fire if years since the last fire were equal to or less than twice the mean reference return interval for a given forest type (e.g., $\leq 2 \times 11$ years for dry mixed conifer) according to the California fire return interval departure

dataset (Safford & Van de Water, 2014). This measure has been used in previous work as an indication of uncharacteristically high fuel accumulation (North et al., 2012) and consequently greater vulnerability to severe fire (Steel et al., 2015). Spatial analyses were performed primarily using the *sf*, *terra*, and *landscape metrics* packages in the R statistical environment (Hesselbarth et al., 2019; Hijmans, 2022; Pebesma, 2018; R Core Team, 2021).

RESULTS

Conifer forest cover decline

As of 2011, there were a total of 1,407,597 hectares of conifer forest in our southern Sierra Nevada study region. 96,810 ha of these forests were classified as either moderate density (41,175 ha) or high density (55,634 ha) mature forest habitat (Table 2). Between 2011 and 2020, canopy cover across 30% of conifer forests declined below 25%, constituting a change to either sparse woodland/savanna vegetation or, in many cases, a transition to nontree-dominated vegetation (Figures 2 and 3). During this period, 50% of moderate or high density mature forest habitat saw canopy cover decline below 40%, constituting a transition to lower density forest (22% of the original extent) or nonforest vegetation (28% of the original extent). Within the mature forest classification, higher density areas experienced more extensive declines, with 85% of this subclass falling below the 60% canopy cover definition of high density. The moderate density mature forests experienced little net change, with much of its declining area compensated by transitions from the high density mature forest group (Table 2; Figure 3).

Spotted owl PACs saw rates of decline across conifer forest classes similar to the study area overall, with the exception of moderate density mature forests, which saw a net increase of 13% due to transitions from the high density class (Table 2). However, because of how they were delineated, PACs generally were biased toward containing denser forests than the study area as a whole, resulting in differences in the average rates of canopy cover decline. In 2011, PACs were characterized by a mean canopy cover of 56%, mean large-tree height of 24 m, and 20% of their conifer forest area were categorized as mature forest of moderate or high density. In contrast, all conifer forests in our study area in 2011 were characterized by a mean canopy cover of 36%, mean large-tree height of 13 m, and 7% were categorized as mature forest of moderate or high density. By 2020, the mean absolute decline in canopy cover was greater inside PACs (26%) compared with all conifer forests in the study area (16%).

Statistical comparisons of PAC and non-PAC areas within Forest Service lands confirmed that the decline in relative canopy cover (percent of 2011 canopy cover) was greater in PACs and also showed clear effects of initial forest structure, with greater decline within areas characterized by higher 2011 canopy cover and less decline in areas characterized by greater 2011 large-tree height (Figure 4). Models predict the average PAC lost 49% (95% prediction interval [PI] = 46, 51) of its 2011 canopy cover by 2020, whereas non-PAC areas lost 42% (PI = 40, 44; Figure 4a). In both PAC and non-PAC areas, areas that had higher canopy cover in 2011 were more susceptible to percentage canopy cover loss and vegetation class transition than areas with lower canopy cover (Figure 4b). For example, mean predicted declines at our density class

TABLE 2 Hectares (percentage of 2011 area) of conifer forests and forest class change between 2011 and 2020 for the full study area and spotted owl protected activity centers (PACs).

Extent	Class	2011	2020	Transition to nonforest	Net change
Study area	All forests	1,407,597	988,601 (70%)	418,996 (30%)	-418,996 (-30%)
	Mature forests ^a	96,810	48,385 (50%)	26,675 (28%)	-48,425 (-50%)
	Mod density	41,175	39,957 (96%)	11,409 (28%)	-1218 (-3%)
	High density	55,634	8427 (15%)	15,265 (27%)	-47,207 (-85%)
Protected activity centers	All forests	76,819	49,936 (65%)	26,883 (35%)	-26,883 (-35%)
	Mature forests ^a	15,072	7568 (50%)	4194 (28%)	-7504 (-50%)
	Mod density	5233	5889 (113%)	1690 (32%)	655 (13%)
	High density	9838	1679 (17%)	2503 (25%)	-8159 (-83%)

Note: Conifer-dominated areas with >25% live canopy cover are considered forests. Within that class, mature forests are defined by a mean tall tree height of >30 m further nested within the mature forest class are moderate density (40%–60% canopy cover) and high density (>60% canopy cover) mature forests. Forests density classes either transitioned to nonforest (canopy cover dropped below 25%) or to a lower density class (e.g., a decline from 70% to 50% cover constitutes a transition from high density mature forest to the moderate density class).

^aIncludes mature forests with >40% 2011 canopy cover (i.e., excludes low density mature forests).




FIGURE 3 Change in conifer forest habitat between 2011 and 2020 in the southern Sierra Nevada. Annual density distribution of canopy cover for (a) all conifer forests (>25% canopy cover), and (b) moderate and high density mature conifer forests (>40% canopy cover and >30 m mean height of tallest trees). Gray portions of density distributions illustrate the area of transition since 2011 of each forest class. (c) Change in area of all conifer forests and mature conifer forests relative to 2011. Mature forest subclasses are shown as dashed lines in (c) and illustrate the initial increase in moderate density mature forests (40%–60% canopy cover) due to transitions from declining high density mature forests (>60% cover).

thresholds of 25%, 40% and 60% canopy cover were 31% (PI = 27, 34), 39% (PI = 37, 41), and 51% (PI = 49, 53), respectively (Figure 4b). The effect of large-tree height on relative canopy cover decline was negative, albeit with a smaller effect size, indicating the presence of large, tall trees had a moderating effect on disturbance. For example, a forest with a mean large-tree height of 15 m was

predicted to experience a 55% (PI = 49, 61) relative canopy cover decline, whereas a mean large-tree height of 30 m was expected to result in a relative canopy cover decline of 52% (PI = 47, 57; Figure 4c) when controlling for 2011 canopy cover. Model uncertainty was low (95% credible intervals did not include zero) for all effect estimates (Appendix S1: Table S1).




FIGURE 4 Expected relative canopy cover declines between 2011 and 2020 as predicted by (a) spotted owl protected activity center (PAC) class, (b) 2011 canopy cover, and (c) 2011 mean large-tree height. Relative canopy cover declines are calculated as the percentage of 2011 canopy cover that was lost between 2011 and 2020. The effects of 2011 canopy cover and height are fitted using data from both PAC and non-PAC sample areas. Points and vertical lines represent the mean estimate and 90% credible intervals in (a). Black lines and grey bands represent the mean estimate and 90% credible intervals in (b) and (c). Tabulated model coefficients can be found in Appendix S1: Table S1.

Wildfire and drought patterns

Because drought conditions were ubiquitous during large parts of our study period, we compared three types of interacting forest disturbances: (1) drought and associated beetle infestation (jointly referred to as “drought”) in the absence of other disturbances, (2) the combination of wildfire plus drought, and (3) the combination of mechanical management activities (e.g., timber harvest and thinning) plus drought. A greater cumulative area of forests was converted to nonforest due to drought alone (213,000 ha; 51% of total area transitioned) than when wildfires were also a contributing factor (190,000 ha; 45%), or when mechanical activities were a contributing factor (16,000 ha; 4%). The differential was even greater in the case of moderate and dense mature forests where transitions to nonforest or a lower density class were more attributable to drought alone (32,000 ha; 66%), than to the combination of wildfire and drought (15,000 ha; 31%) or mechanical activities and drought (2000 ha; 4%).

For all forest classes, patch and core size of transitioned areas were greater within burned forest than those that were not burned during 2011–2020 (Figure 2; Table 3). Distributions of disturbed patch sizes were highly skewed with relatively few large patches accounting for a disproportionate amount of the area of transitioned forest. The largest transitioned forest patch (68,389 ha) and contiguous core area (25,186 ha) of transitioned forest were found within burned areas during

this period. Likewise, the largest patch size (1119 ha) and contiguous core area (326 ha) of transitioned moderate and high density mature forest habitat occurred within fire perimeters (Table 3). These differences in spatial pattern of interacting drought and wildfire resulted in greater forest fragmentation than areas affected by drought alone as measured by the aggregation index. Low aggregation index values approaching 0 represent greater fragmentation and high values approaching 100 represent greater aggregation. For all forests, the aggregation index declined from 93 in 2011 to 88 in unburned areas and to 75 in burned areas by 2020. Similarly, the aggregation index for mature forests declined from 85 in 2011 to 75 in unburned forests and 64 in burned forests (Table 4).

There was a clear temporal pattern of forest mortality attributable to drought and beetle infestation alone with a 2016 peak in forest cover decline corresponding to the last year of the 2012–2016 drought. Transitions due to the combination of wildfire and drought were less consistent and corresponded with individual large wildfire events (e.g., the 2013 Rim Fire and the 2020 Creek and Castle Fires). Transitions attributable to mechanical activities and drought were consistently limited between 2012 and 2019 (Figure 5a).

As of 2011, a large majority of the conifer forests and mature forests in the southern Sierra Nevada had not experienced a wildfire in at least twice their reference fire return interval. The absolute amount of forest

TABLE 3 Summary statistics of disturbance patch size and core area size created by drought/beetles and wildfire between 2011 and 2020.

Metric	Class	Disturbance	N	Mean	Q50	Q95	Max
Area	Forest	Drought	233,874	0.66	0.18	1.89	2246.04
Area	Forest	Fire + Drought	36,953	7.15	0.18	3.77	68,372.36
Area	High density mature	Drought	11,467	2.25	0.27	5.84	1741.95
Area	High density mature	Fire + Drought	5694	3.74	0.36	9.52	827.43
Area	Mature forest	Drought	20,306	1.04	0.18	3.23	447.88
Area	Mature forest	Fire + Drought	9078	3.00	0.36	8.17	1115.57
Area	Moderate density mature	Drought	14,629	0.86	0.18	2.51	373.26
Area	Moderate density mature	Fire + Drought	6939	1.54	0.27	4.49	970.91
Core	Forest	Drought	1268	2.36	0.54	7.54	342.46
Core	Forest	Fire + Drought	671	116.43	0.81	93.79	25,179.57
Core	High density mature	Drought	375	6.15	0.63	24.90	653.32
Core	High density mature	Fire + Drought	322	10.15	1.39	41.56	295.41
Core	Mature forest	Drought	278	2.51	0.36	11.57	67.88
Core	Mature forest	Fire + Drought	400	8.41	0.99	32.24	326.03
Core	Moderate density mature	Drought	124	2.68	0.45	10.98	58.63
Core	Moderate density mature	Fire + Drought	156	5.40	0.72	19.64	292.45

Note: Patches without core area are included in the patch size (area) metric but excluded from the core area (core) metric. *N* represents the number of patches in each group, and all other summary statistics are in hectares. In most cases disturbed patches are small (i.e., 50th quantile <1 ha). Thus, we summarize the large end of data distributions (i.e., 95th quantile, and max) as these patches affect the greatest amount of area and are the most ecologically important.

TABLE 4 Aggregation index values prior to and following drought/beetles and wildfire in the southern Sierra Nevada during 2011–2020.

Class	Initial (2011)	Postdrought (2020)	Postdrought and wildfire (2020)
All forests	93	88	75
Mature forests	85	75	64
Moderate density mature	77	63	52
High density mature	83	61	46

Note: Index values range from 0 to 100, in which low values represent highly fragmented forest cover and high values represent highly aggregated forest cover.

that was within twice its reference return interval was largely stable across the study period, but the proportion of recently burned forest and recently burned mature forest increased due to an overall decline in forest cover (Figure 5b). Specifically, the percentage of recently burned conifer forests increased from 25% in 2011 to 33% in 2020, whereas recently burned mature forests increased from 7% to 20% during the same period.

DISCUSSION

Rapid decline of mature forest habitat

The 2010s was a decade characterized by severe and unprecedented ecological disturbance in California's conifer forests; one that portends a rapid reshaping of forested landscapes, the ecosystem services they provide, and their dependent wildlife communities. Here we documented a temporary or permanent conversion of 30% of the conifer forests and a transition of half of the moderate and high density mature forest habitat in the southern Sierra Nevada between 2011 and 2020 due to compounding megafires and a historic drought. Changes were particularly pronounced in dense mature forest habitat, which declined by 85% either by transitioning to lower density forest or through complete conversion to nonforest vegetation. Spotted owl PACs, also saw dramatic declines with a loss of 49% of their 2011 canopy cover. Worsening wildfire and climate trends are expected to continue or accelerate in the coming decades (Abatzoglou et al., 2021), making the continued decline of conifer forest extent likely and the complete loss of remaining moderate and high density mature forest habitats plausible (Stephens, Miller, et al., 2016; Westerling, 2016).




FIGURE 5 (a) All forest and mature forest area transitioned annually due to drought, a combination of drought and wildfire, or a combination of drought and mechanical management activities (timber harvest or thinning) from 2011 to 2020. (b) The total amount of area remaining as forest or mature forest as well as the amount of area that survived a recent burn (time since the last fire was less than twice the historic fire return interval). Conifer forests are defined by a minimum of 25% canopy cover. Mature forests are defined as having a minimum of 30 m average height of the tallest trees as of 2011 and maintaining at least 40% canopy cover. The timing of three large fires and the final year of the 2012–2016 drought are noted with vertical dashed lines. No mechanical activity data were available during 2020, potentially inflating the estimate of drought-only transitions for that year. Changes in mature forest density subclasses are shown in Appendix S1: Figure S1.

Megafires create large high severity patches that can be inaccessible or inhospitable to mature forest specialists, and as a result they tend to support lower wildlife species diversity overall (Jones et al., 2020; Steel, Fogg, et al., 2022). Further, large severe wildfires can increase fragmentation of limited mature forest habitat. Our results demonstrate that wildfire (in combination with drought) created distinctly larger contiguous patches of forest loss and increased forest fragmentation, whereas tree mortality from drought and beetle kill resulted in patchier and more fine-scale spatial patterns with relatively moderate increases in habitat fragmentation. In the Sierra Nevada, suitable fisher habitat is distributed in a north/south pattern limited by elevational bounds and periodically disrupted by bottlenecks that occur near large river canyons (Thompson et al., 2021). These bottlenecks, which are typically associated with isolated stretches of linkage habitat, may be comprised of lower quality habitat but are crucial for maintaining overall population connectivity and gene flow (Tucker et al., 2014). Since 2013, habitat within key linkage areas has been lost to high severity fire at nearly twice the rate of habitat outside key linkage areas (Thompson et al., 2021; USFWS, 2020), limiting population-scale connectivity and increasing the risk of genetic isolation (Tucker et al., 2014). The spotted owl is tolerant of, or even selects for, small high severity burned patches if mature stands survive in the nearby matrix, but they are intolerant of increasingly large high severity patches created by fires such as the 2013 Rim, 2020 Creek, and 2020 Castle fires in the southern Sierra Nevada (Jones et al., 2020; Kramer et al., 2021). A mosaic of burn severities consisting of predominantly low-to-moderate severity fire with small patches of high severity can create “pyrodiverse” landscapes that contain early seral habitat including shrublands and recently killed “snag forests” required by some species, while also promoting the persistence of key habitat structures for late seral-dependent species (Jones, Kramer, et al., 2021; Steel, Collins, et al., 2021). These disturbances also leave a network of residual habitat patches (i.e., fire refugia) that are critical to fisher re-colonization (Blomdahl, 2018; Thompson et al., 2021). Without these refugia, the potential exists for a community shift to more generalist species (Green et al., 2022).

Low-to-moderate severity fire can create habitat for some wildlife species and increase forest resistance to future severe disturbance (Steel, Foster, et al., 2021), but drought mortality has the potential to add flammable fuels to the system and increase the likelihood of future forest loss due to wildfire (Stephens et al., 2018, 2022). The 2012–2016 drought and associated beetle epidemics resulted in the mortality of more than 150 million trees in California’s Sierra Nevada (USDA, 2020), which was

disproportionately concentrated in large conifers (e.g., in >38 cm diameter ponderosa pine; Fettig et al., 2019). Fishers rely on large trees for resting and denning habitat (Green et al., 2019; Purcell et al., 2009), and spotted owl population declines have been linked to losses of large, old trees used for nesting (Jones et al., 2018). Thus, the mechanism of canopy cover decline in mature forests is also important. For example, shifts from high density to moderate density mature forests may be restorative when wildfire is the causal agent because it consumes fuel and generally kills smaller trees, leaving larger trees intact and further insulated from subsequent disturbance. Fuels reduction treatments that target fine fuels can have a similarly restorative effect, especially when including the use of prescribed or cultural fire (North et al., 2021). Additional research is needed to better understand how mature forest specialist species respond to wildfire and wildfire surrogates that reduce canopy cover below a known optimum, while maintaining important features such as large, tall conifers and restoring resilience to future disturbance (Jones, Keyser, et al., 2021). However, when canopy cover declines are attributable to drought and beetle attack (Appendix S1: Figure S1), there is likely to be a corrosive effect on habitat quality because large trees are preferentially killed and dead fuel is added to the system, increasing the vulnerability to subsequent wildfire effects.

Rapid loss in mature forest habitat in the southern Sierra Nevada and longer term trends in fire-related forest decline throughout California (Steel et al., 2018; Stevens et al., 2017) suggest that existing forest management paradigms may be inadequate for maintaining mature mixed-conifer forests under current and projected future disturbance dynamics (North et al., 2022). If these rates of decline continue, we are likely to see near total loss of southern Sierra Nevada mature conifer forests in the coming decades. This would be much more rapid than the time horizon of mature forest loss estimated by Stephens and colleagues (2016) (by 2089, or ~75 years). However, Stephens and colleagues (2016) did not consider drought-related mortality, and only analyzed fire activity up to 2014, which missed the record fire year of 2020 (Safford et al., 2022). It is worth noting that the extreme fire activity documented in California during 2020 was likely not a one-off anomaly; recent observations indicate similar, if not exacerbated, fire activity in 2021 (Shive et al., 2021). The region has also re-entered extreme drought (Williams et al., 2022) with implications for both drought and beetle mortality and severe wildfire. More optimistically, total loss of mature forests in this region could be delayed until midcentury if we enter a period of cooler, wetter years, if surviving mature forests within these fire footprints have gained resilience to future disturbances, or if recruitment

of mature hardwood species compensates for losses of large conifers. Hardwood species may become a greater component of the Sierra Nevada landscape as conifers decline (Restaino et al., 2019; Steel, Foster, et al., 2021). Oaks, especially California black oaks, are relatively resilient to both wildfire and drought, and are utilized by species such as the spotted owl and fisher (Aubry et al., 2013; Green et al., 2019; North et al., 2000). However, loss of mature forest habitat, on any likely timeline, is unsustainable given that the recruitment of conifer or hardwood mature forests takes many decades to centuries. Stephens et al. (2016) emphasized that policies prioritizing forest resilience over other resource concerns may be needed to meaningfully address the current backlog in forest management and shift the course from forest decline to sustainable disturbance dynamics. Indeed, our analysis showed that areas of higher canopy cover are more at risk of loss, and that large areas of relatively homogenous moderate and higher density forests, like PACs, are at risk of larger declines if resilience needs are not addressed. Recognizing the dynamic nature of habitat in these forests, and prioritizing the restoration of these dynamics over the attempted strict preservation of existing habitat, may help to minimize the impacts of these changes and maintain habitat functionality in the long term (Fabritius et al., 2017; Gaines et al., 2022; Stoetzel et al., 2020).

Management implications: Toward more sustainable disturbance dynamics

Landscapes are naturally dynamic in space and time, but society's approach for conserving landscapes often views them as static entities (Stoetzel et al., 2020). Habitat conservation typically involves cordoning off lands for full or partial protection from human activities, while at the same time allowing for fire suppression activities to preserve current structure. Conservation approaches that aim to keep ecosystems in stasis have been largely successful in conserving biodiversity and threatened species worldwide (Gray et al., 2016) by preventing threats to habitats such as infrastructure development and natural resource extraction (Barber et al., 2014). These approaches are heavily informed by conservation objectives in more naturally static ecosystems (e.g., Barber et al., 2014) as opposed to those adapted to frequent low intensity disturbance. Such "static" conservation approaches are heavily embedded in existing wildlife and ecosystem conservation policy (Leopold et al., 2018), as well as land management plans (e.g., USDA, 2004) in North America. Yet recent disturbance patterns and their cumulative impacts have demonstrated that efforts to resist change are often falling short in

dynamic ecosystems, such that achieving the specific conservation objectives and possibly the intent outlined in policy documents may no longer be feasible in disturbance-prone areas (Davis et al., *In Press*). In fact, continued attempts to resist change may be counterproductive when a hands-off approach (but continued fire suppression) creates a higher likelihood of rapid, transformational, and undesirable changes in the form of large-scale type conversion and habitat loss from disturbance (Rissman et al., 2018). In our study region, spotted owl PACs are often managed using a static conservation approach but our analysis shows that they have recently experienced more declines in canopy cover (49% relative to 2011) than outside of their borders (4%). This observation suggests that conservation of habitat for old-forest-dependent species may require a more dynamic approach that increases the resilience to disturbance, while maintaining valuable habitat features such as large, tall trees.

As an alternative to the static approach to habitat conservation currently in practice, a landscape conservation paradigm that recognizes and incorporates ecological system dynamics (Gaines et al., 2022; Hessburg et al., 2021) may prove to be better suited for disturbance-prone forests. Collectively, we refer to this as the "managed dynamics" paradigm (Figures 1 and 6). Although the overall goal of this alternative approach is similar to that of a more traditional static paradigm, i.e., to conserve a particular landscape feature (in this case, mature forests), the managed dynamics paradigm strives for something closer to a dynamic equilibrium (*sensu* Bonnicksen & Stone, 1982) of habitat loss and recruitment. In Sierra Nevada dry forests that historically were highly dynamic, patches of mature forest were periodically lost or degraded but were balanced by continual successional changes supported by a frequent low severity disturbance regime (Miller & Safford, 2017). Moving away from a "static" conservation paradigm in favor of a "dynamic" one does not prescribe eliminating protected areas or habitat preserves; nor would it involve removal of large trees, which our analysis shows supported forest resilience during the last decade. Rather it suggests that greater active management is necessary (e.g., through fire use and ecologically based thinning; *sensu* North et al., 2021) with an eye for emulating fine-scale heterogeneity and forest resilience likely supported by historic disturbance regimes (Jones, Keyser, et al., 2021). Further, management tactics and tools may need to be similarly dynamic and allowed to shift when objectives are not being met, or when needs have changed (e.g., Spies et al., 2018; Wood et al., 2018).

Managing for sustainable dynamics also embraces the positive role of humans in actively managing ecosystems. For millennia, Indigenous peoples of western North America burned vast areas of land to increase access to natural




FIGURE 6 Illustration of two approaches to management in seasonally dry disturbance-prone California forests. Attempts to avoid and exclude all disturbance has led to mega-disturbances and transitions from mature forests conditions (top scenario), while managing for sustainable disturbance dynamics can maintain forest resilience and wildlife habitat (bottom scenario).

resources (Anderson, 2013; Hoffman et al., 2021). Indigenous burning as part of historic fire regimes also produced other ecosystem benefits such as increased forest resilience to subsequent disturbances (Eisenberg et al., 2019), enhanced biodiversity (Hoffman et al., 2021), and mutual protection for cultural and ecological resources (Slaton et al., 2019). A major part of implementing a managed dynamics paradigm in the Sierra Nevada includes increasing support for Indigenous fire use, as well as state and federally supported and ecologically driven management. Specific management actions would likely include leveraging areas of low-to-moderate severity effects within unplanned wildfires as well as active fuels reductions across large landscapes with prescribed and managed fire in a “pyrosilviculture” framework (North et al., 2021). Such an approach would be expected to result in reduced forest conversion to nonforest, increased water supply, more stable carbon storage, reduced competitive stress, as well as both protection and promotion of growing conditions for large, old trees (Figure 6; Stephens et al., 2020, 2021; North et al., 2022).

A managed dynamics paradigm is not without its challenges. Implementing forest treatments and restorative actions is generally difficult due to insufficient funding

and staffing, perceived and realized risks of employing prescribed, cultural, and managed wildfire, as well as other political and societal concerns (North et al., 2021; Schultz et al., 2019). Perhaps foremost among these obstacles when sensitive wildlife habitat is of concern is that a managed dynamics approach appears at odds with the precautionary principle, which has governed land management and wildlife conservation for decades (e.g., Kriebel et al., 2001). The precautionary principle “imposes a burden of proof on those who create potential risks, and it requires the regulation of activities, even if it cannot be shown that those activities are likely to produce significant harms” (Sunstein, 2002). In the Sierra Nevada, the principle has led to resistance of managing disturbance dynamics because management activities such as fuels reduction and thinning have been perceived to pose a risk to spotted owls and fishers via habitat alteration (Tempel et al., 2015). Indeed, many past management practices have had negative impacts on old-forest habitat features, including harvesting of the largest trees and removal of damaged trees that contain cavities or provide platforms for nesting and denning. Yet under current conditions characterized by rapid change, decision “paralysis”

(Sunstein, 2002) precipitated by the precautionary principle may lead to far greater harm to spotted owls and fishers via complete loss of mature forest habitat (Hessburg et al., 2021). For example, an estimated 10%–14% of the global population of giant sequoia were lost in 2020 alone due to high severity wildfire within the southern Sierra Nevada with additional losses to giant sequoia groves incurred from two large wildfires in 2021 (Shive et al., 2021). Many of the affected giant sequoia groves had largely gone untreated (i.e., no large-scale fuels reduction activities have occurred) and there is concern that increased restrictions for the southern Sierra fisher population may add an additional barrier to restoration efforts. Yet avoidance of action is a management decision, and our analysis shows that a hands-off approach is increasingly failing to preserve mature forests. Thus, we suggest that the managed dynamics paradigm is compatible with an interpretation of the precautionary principle that explicitly recognizes the risk of inaction. For example, as described in Wood et al. (2020), the precautionary principle could posit that management actions should be taken despite uncertainties if the cost of inaction is high.

CONCLUSION

Shifting disturbance patterns pose stark challenges to forest management and conservation. This has become particularly evident during the past decade in the southern Sierra Nevada of California, where compounding megafires and unprecedented drought have rapidly eroded already limited mature forest habitat. Although this period of forest degradation appears exceptional at present, continued climate change and the lingering legacy of fire exclusion and suppression make continued mega-disturbances likely. Indeed, drought conditions returned to the region in 2019 and persist through the time of writing in 2022. To forestall continued loss and fragmentation of habitat critical to mature forest specialist species a shift from a static, preservationist paradigm of habitat conservation to one striving for sustainable disturbance dynamics is likely to be necessary in fire- and drought-prone forests. Encouragingly, some recent forest policy appears to be moving toward a recognition of the need for managing for desirable dynamics. For example, recent management strategies for both Sierra Nevada fishers and California spotted owls recognized the shortcomings of the static approach of identifying “protected” habitat areas with limited management allowed (USDA, 2019). These new strategies call for a more dynamic approach to providing this habitat over time and space, while also incorporating the likelihood of disturbance.

A broadscale shift in forest conservation paradigm faces challenges, but given the rapid rate of forest degradation and future outlook, an equally rapid adaptation of management approach is likely to be needed to maintain and restore mature forests in the coming decades.

ACKNOWLEDGMENTS

We thank the USFS Region 5 remote-sensing laboratory for providing the F^3 data used in this analysis, especially Laura Young-Hart and Leland Tarnay. Figure 6 was illustrated by the artist Allison Fitzmorris. Funding was provided by the California Department of Forestry and Fire Protection, grant #8GG21823.

CONFLICT OF INTEREST

The authors have no conflicts of interest.

DATA AVAILABILITY STATEMENT

Code (Steel, Jones, et al., 2022a) is available in Zenodo at <https://doi.org/10.5281/zenodo.7044555>. Data (Steel, Jones, et al., 2022b) are available in Dryad at <https://doi.org/10.6078/D1W43V>. Additional publicly available data sets utilized for this research are as follows: California fire perimeter geospatial data (FRAP, 2022; <https://frap.fire.ca.gov/frap-projects/fire-perimeters/>) and the California fire return interval departure dataset (Safford & Van de Water, 2014; <https://www.fs.usda.gov/detail/r5/landmanagement/gis/?cid=stelprdb5327836>).

ORCID

Zachary L. Steel  <https://orcid.org/0000-0002-1659-3141>

Gavin M. Jones  <https://orcid.org/0000-0002-5102-1229>

REFERENCES

- Abatzoglou, J. T., D. S. Battisti, A. P. Williams, W. D. Hansen, B. J. Harvey, and C. A. Kolden. 2021. “Projected Increases in Western US Forest Fire despite Growing Fuel Constraints.” *Communications Earth & Environment* 2: 1–8.
- Anderson, M. K. 2013. *Tending the Wild: Native American Knowledge and the Management of California's Natural Resources*. Berkeley, CA: University of California Press.
- Aubry, K. B., C. M. Raley, S. W. Buskirk, W. J. Zielinski, M. K. Schwartz, R. T. Golightly, K. L. Purcell, R. D. Weir, and J. S. Yaeger. 2013. “Meta-Analyses of Habitat Selection by Fishers at Resting Sites in the Pacific Coastal Region.” *The Journal of Wildlife Management* 77: 965–74.
- Barber, C. P., M. A. Cochrane, C. M. Souza, and W. F. Laurance. 2014. “Roads, Deforestation, and the Mitigating Effect of Protected Areas in the Amazon.” *Biological Conservation* 177: 203–9.
- Bernal, A. A., S. L. Stephens, B. M. Collins, and J. J. Battles. 2022. “Biomass Stocks in California's Fire-Prone Forests: Mismatch in Ecology and Policy.” *Environmental Research Letters* 17: 044047.

- Blomdahl, E. M. 2018. "Interactions Between Fire Severity and Forest Biota in the Central Sierra Nevada: Formation and Impact of Small-scale Fire Refugia and the Effect of Fire on Forest Structure Predictive of Fisher (*Pekania pennanti*) Den Habitat." MS thesis. Logan, UT: Utah State University.
- Bonnicksen, T. M., and E. C. Stone. 1982. "Reconstruction of a Presettlement Giant Sequoia-Mixed Conifer Forest Community Using the Aggregation Approach." *Ecology* 63: 1134–48.
- Brewer, C., B. Schwind, R. Warbington, W. Clerke, P. Krosse, L. Suring, and M. Schanta. 2005. "Section 3: Existing Vegetation Mapping Protocol." In *Existing Vegetation Classification and Mapping Technical Guide*. Gen. Tech. Rep. WO-67, edited by R. Brohman and L. Bryant. Washington, DC: USDA-Forest Service, Ecosystem Management Coordination Staff.
- Bürkner, P.-C. 2017. "Brms: An R Package for Bayesian Multilevel Models Using Stan." *Journal of Statistical Software* 80: 1–28.
- Collins, B. M., D. L. Fry, J. M. Lydersen, R. Everett, and S. L. Stephens. 2017. "Impacts of Different Land Management Histories on Forest Change." *Ecological Applications* 27: 2475–86.
- Collins, B. M., S. L. Stephens, J. J. Moghaddas, and J. Battles. 2010. "Challenges and Approaches in Planning Fuel Treatments across Fire-Excluded Forested Landscapes." *Journal of Forestry* 108: 24–31.
- Cunningham, S. C., W. B. Ballard, L. M. Monroe, M. J. Rabe, and K. D. Bristow. 2003. "Black Bear Habitat Use in Burned and Unburned Areas, Central Arizona." *Wildlife Society Bulletin (1973-2006)* 31: 786–92.
- Davis, R. J., D. M. Bell, M. J. Gregory, Z. Yang, A. N. Gray, and A. E. Stratton. In Press. "Northwest Forest Plan – the First 25 Years: Status and Trends of Late-Successional and Old-Growth Forests." Gen. Tech. Rep. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station.
- Eisenberg, C., C. L. Anderson, A. Collingwood, R. Sissons, C. J. Dunn, G. W. Meigs, D. E. Hibbs, et al. 2019. "Out of the Ashes: Ecological Resilience to Extreme Wildfire, Prescribed Burns, and Indigenous Burning in Ecosystems." *Frontiers in Ecology and Evolution* 7: 436.
- Fabritius, H., A. Jokinen, and M. Cabeza. 2017. "Metapopulation Perspective to Institutional Fit: Maintenance of Dynamic Habitat Networks." *Ecology and Society* 22: 30.
- Fettig, C. J., L. A. Mortenson, B. M. Bulaon, and P. B. Foulk. 2019. "Tree Mortality Following Drought in the Central and Southern Sierra Nevada, California, U.S." *Forest Ecology and Management* 432: 164–78.
- FRAP. 2022. *Fire and Resource Assessment Program Fire Perimeter Data*. Sacramento, CA: California Department of Forestry and Fire Protection. <https://frap.fire.ca.gov/frap-projects/fire-perimeters/>.
- Gaines, W. L., P. F. Hessburg, G. H. Aplet, P. Henson, S. J. Prichard, D. J. Churchill, G. M. Jones, D. J. Isaak, and C. Vynne. 2022. "Climate Change and Forest Management on Federal Lands in the Pacific Northwest, USA: Managing for Dynamic Landscapes." *Forest Ecology and Management* 504: 119794.
- Gray, C. L., S. L. Hill, T. Newbold, L. N. Hudson, L. Börger, S. Contu, A. J. Hoskins, S. Ferrier, A. Purvis, and J. P. W. Scharlemann. 2016. "Local Biodiversity Is Higher inside than outside Terrestrial Protected Areas Worldwide." *Nature Communications* 7: 12306.
- Green, D. S., M. E. Martin, R. A. Powell, E. L. McGregor, M. W. Gabriel, K. L. Pilgrim, M. K. Schwartz, and S. M. Matthews. 2022. "Mixed-Severity Wildfire and Salvage Logging Affect the Populations of a Forest-Dependent Carnivoran and a Competitor." *Ecosphere* 13: e03877.
- Green, R. E., K. L. Purcell, C. M. Thompson, D. A. Kelt, and H. U. Wittmer. 2019. "Microsites and Structures Used by Fishers (*Pekania Pennanti*) in the Southern Sierra Nevada: A Comparison of Forest Elements Used for Daily Resting Relative to Reproduction." *Forest Ecology and Management* 440: 131–46.
- Gutiérrez, R. J., P. N. Manley, and P. A. Stine. 2017. "The California spotted owl: current state of knowledge." Gen. Tech. Rep. PSW-GTR-254. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station 254.
- Hessburg, P. F., D. J. Churchill, A. J. Larson, R. D. Haugo, C. Miller, T. A. Spies, M. P. North, et al. 2015. "Restoring Fire-Prone Inland Pacific Landscapes: Seven Core Principles." *Landscape Ecology* 30: 1805–35.
- Hessburg, P. F., S. J. Prichard, R. K. Hagmann, N. A. Povak, and F. K. Lake. 2021. "Wildfire and Climate Change Adaptation of Western North American Forests: A Case for Intentional Management." *Ecological Applications* 31: e02432.
- Hesselbarth, M. H. K., M. Sciaini, K. A. With, K. Wiegand, and J. Nowosad. 2019. "LandscapeMetrics: An Open-Source R Tool to Calculate Landscape Metrics." *Ecography* 42: 1648–57.
- Hijmans, R. 2022. "terra: Spatial Data Analysis." <https://rspatial.org/terra/>.
- Hoffman, K. M., E. L. Davis, S. B. Wickham, K. Schang, A. Johnson, T. Larking, P. N. Lauriault, N. Q. Le, E. Swerdfager, and A. J. Trant. 2021. "Conservation of Earth's Biodiversity Is Embedded in Indigenous Fire Stewardship." *Proceedings of the National Academy of Sciences* 118: e2105073118.
- Huang, S., C. Ramirez, M. McElhaney, and K. Evans. 2018. "F3: Simulating Spatiotemporal Forest Change from Field Inventory, Remote Sensing, Growth Modeling, and Management Actions." *Forest Ecology and Management* 415–416: 26–37.
- Hutto, R. L. 2008. "The Ecological Importance of Severe Wildfires: Some like it Hot." *Ecological Applications* 18: 1827–34.
- Jones, G. M., J. J. Keane, R. J. Gutiérrez, and M. Z. Peery. 2018. "Declining Old-Forest Species as a Legacy of Large Trees Lost." *Diversity and Distributions* 24: 341–51.
- Jones, G. M., A. R. Keyser, A. L. Westerling, W. J. Baldwin, J. J. Keane, S. C. Sawyer, J. D. Clare, R. Gutiérrez, and M. Z. Peery. 2021. "Forest Restoration Limits Megafires and Supports Species Conservation under Climate Change." *Frontiers in Ecology and the Environment* 20: 210–6.
- Jones, G. M., H. A. Kramer, W. J. Berigan, S. A. Whitmore, R. J. Gutiérrez, and M. Z. Peery. 2021. "Megafire causes persistent loss of an old-forest species." *Animal Conservation* 24: 925–36.
- Jones, G. M., H. A. Kramer, S. A. Whitmore, W. J. Berigan, D. J. Tempel, C. M. Wood, B. K. Hobart, et al. 2020. "Habitat Selection by Spotted Owls after a Megafire Reflects their Adaptation to Historical Frequent-Fire Regimes." *Landscape Ecology* 35: 1199–213.
- Knapp, E. E., C. N. Skinner, M. P. North, and B. L. Estes. 2013. "Long-Term Overstory and Understory Change Following Logging and Fire Exclusion in a Sierra Nevada Mixed-Conifer Forest." *Forest Ecology and Management* 310: 903–14.

- Knight, C. A., R. E. Tompkins, J. A. Wang, R. York, M. L. Goulden, and J. J. Battles. 2022. "Accurate Tracking of Forest Activity Key to Multi-Jurisdictional Management Goals: A Case Study in California." *Journal of Environmental Management* 302: 114083.
- Koltunov, A., C. M. Ramirez, S. L. Ustin, M. Slaton, and E. Haunreiter. 2020. "eDaRT: The Ecosystem Disturbance and Recovery Tracker System for Monitoring Landscape Disturbances and their Cumulative Effects." *Remote Sensing of Environment* 238: 111482.
- Koltunov, A., C. Ramirez, and S. Ustin. 2015. "eDaRT: the Ecosystem Disturbance and Recovery Tracking system prototype supporting ecosystem management in California." 2015 NASA Carbon Cycle and Ecosystems Joint Science Workshop, College Park, MD.
- Kramer, H. A., G. M. Jones, S. A. Whitmore, J. J. Keane, F. A. Atuo, P. B. Dotters, S. C. Sawyer, S. L. Stock, R. J. Gutiérrez, and M. Z. Peery. 2021. "California Spotted Owl Habitat Selection in a Fire-Managed Landscape Suggests Conservation Benefit of Restoring Historical Fire Regimes." *Forest Ecology and Management* 479: 118576.
- Kriebel, D., J. Tickner, P. Epstein, J. Lemons, R. Levins, E. L. Loechler, M. Quinn, R. Rudel, T. Schettler, and M. Stoto. 2001. "The Precautionary Principle in Environmental Science." *Environmental Health Perspectives* 109: 871–6.
- Leopold, B. D., W. B. Kessler, and J. L. Cummins. 2018. *North American Wildlife Policy and Law*. Missoula, MT: Boone and Crockett Club.
- Liang, S., M. D. Hurteau, and A. L. Westerling. 2018. "Large-Scale Restoration Increases Carbon Stability under Projected Climate and Wildfire Regimes." *Frontiers in Ecology and the Environment* 16: 207–12.
- Lindenmayer, D. B., and W. F. Laurance. 2017. "The Ecology, Distribution, Conservation and Management of Large Old Trees." *Biological Reviews* 92: 1434–58.
- Lindenmayer, D. B., W. F. Laurance, and J. F. Franklin. 2012. "Global Decline in Large Old Trees." *Science* 338: 1305–6.
- Lydersen, J. M., and B. M. Collins. 2018. "Change in Vegetation Patterns over a Large Forested Landscape Based on Historical and Contemporary Aerial Photography." *Ecosystems* 21: 1348–63.
- Mayer, K. E., and W. F. Laudenslayer. 1988. *A Guide to Wildlife Habitats of California*, 166. Sacramento, CA: Department of Fish and Game.
- Meyer, M. D., and M. P. North. 2019. "Natural Range of Variation of Red Fir and Subalpine Forests in the Sierra Nevada Bioregion." Gen. Tech. Rep. U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station, Albany, CA.
- Miller, J. D., E. E. Knapp, C. H. Key, C. N. Skinner, C. J. Isbell, R. M. Creasy, and J. W. Sherlock. 2009. "Calibration and Validation of the Relative Differenced Normalized Burn Ratio (RdNBR) to Three Measures of Fire Severity in the Sierra Nevada and Klamath Mountains, California, USA." *Remote Sensing of Environment* 113: 645–56.
- Miller, J. D., and H. D. Safford. 2017. "Corroborating Evidence of a Pre-Euro-American Low- to Moderate-Severity Fire Regime in Yellow Pine-Mixed Conifer Forests of the Sierra Nevada, California, USA." *Fire Ecology* 13: 58–90.
- North, M. P., B. M. Collins, H. D. Safford, and N. L. Stephenson. 2016. "Montane Forests." In *Ecosystems of California*, edited by H. Mooney and E. Zavaleta, 553–77. Berkeley, CA: University of California Press.
- North, M. P., B. M. Collins, and S. Stephens. 2012. "Using Fire to Increase the Scale, Benefits, and Future Maintenance of Fuels Treatments." *Journal of Forestry* 110: 392–401.
- North, M. P., J. T. Kane, V. R. Kane, G. P. Asner, W. Berigan, D. J. Churchill, S. Conway, et al. 2017. "Cover of tall trees best predicts California spotted owl habitat." *Forest Ecology and Management* 405: 166–78.
- North, M. P., G. Steger, R. Denton, G. Eberlein, T. Munton, and K. Johnson. 2000. "Association of Weather and Nest-Site Structure with Reproductive Success in California Spotted Owls." *Journal of Wildlife Management* 64(3): 797–807.
- North, M. P., R. E. Tompkins, A. A. Bernal, B. M. Collins, S. L. Stephens, and R. A. York. 2022. "Operational Resilience in Western US Frequent-Fire Forests." *Forest Ecology and Management* 507: 120004.
- North, M. P., R. A. York, B. M. Collins, M. D. Hurteau, G. M. Jones, E. E. Knapp, L. Kobziar, et al. 2021. "Pyrosilviculture Needed for Landscape Resilience of Dry Western United States Forests." *Journal of Forestry* 119: 520–44.
- Odion, D. C., and C. T. Hanson. 2013. "Projecting Impacts of Fire Management on a Biodiversity Indicator in the Sierra Nevada and Cascades, USA: The Black-Backed Woodpecker." *The Open Forest Science Journal* 6: 14–23.
- Pebesma, E. 2018. "Simple Features for R: Standardized Support for Spatial Vector Data." *The R Journal* 10: 439–46.
- Purcell, K. L., A. K. Mazzoni, S. R. Mori, and B. B. Boroski. 2009. "Resting Structures and Resting Habitat of Fishers in the Southern Sierra Nevada, California." *Forest Ecology and Management* 258: 2696–706.
- R Core Team. 2021. *R: A Language and Environment for Statistical Computing*. Vienna: R Foundation for Statistical Computing.
- Restaino, C., D. J. N. Young, B. Estes, S. Gross, A. Wuenschel, M. Meyer, and H. Safford. 2019. "Forest Structure and Climate Mediate Drought-Induced Tree Mortality in Forests of the Sierra Nevada, USA." *Ecological Applications* 29: e01902.
- Rissman, A. R., K. D. Burke, H. A. C. Kramer, V. C. Radeloff, P. R. Schilke, O. A. Selles, R. H. Toczylowski, et al. 2018. "Forest Management for Novelty, Persistence, and Restoration Influenced by Policy and Society." *Frontiers in Ecology and the Environment* 16: 454–62.
- Robeson, S. M. 2015. "Revisiting the Recent California Drought as an Extreme Value." *Geophysical Research Letters* 42: 6771–9.
- Safford, H. D., A. K. Paulson, Z. L. Steel, D. J. N. Young, and R. B. Wayman. 2022. "The 2020 California Fire Season: A Year like no Other, a Return to the Past or a Harbinger of the Future?" *Global Ecology and Biogeography* 31: 2005–25.
- Safford, H. D., and J. T. Stevens. 2017. "Natural Range of Variation for Yellow Pine and Mixed-Conifer Forests in the Sierra Nevada, Southern Cascades, and Modoc and Inyo National Forests, California, USA." Gen. Tech. Rep. Albany, CA: U. S. Department of Agriculture, Forest Service, Pacific Southwest Research Station.
- Safford, H. D., and K. M. Van de Water. 2014. "Using Fire Return Interval Departure (FRID) Analysis to Map Spatial and Temporal Changes in Fire Frequency on National Forest Lands in California." Res. Pap. PSW-RP-266. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. 59 p.
- Schultz, C. A., S. M. McCaffrey, H. R. Huber-Stearns, C. A. Schultz, S. M. McCaffrey, and H. R. Huber-Stearns. 2019. "Policy

- Barriers and Opportunities for Prescribed Fire Application in the Western United States.” *International Journal of Wildland Fire* 28: 874–84.
- Shive, K., C. Brigham, T. Caprio, and P. Hardwick. 2021. *2021 Fire Season Impacts on Giant Sequoias*. Three Rivers, CA: National Park Service, Sequoia and Kings Canyon National Parks.
- Slaton, M. R., J. G. Holmquist, M. Meyer, R. Andrews, and J. Beidl. 2019. “Traditional Ecological Knowledge Used in Forest Restoration Benefits Natural and Cultural Resources: The Intersection between Pandora Moths, Jeffrey Pine, People, and Fire.” *Natural Areas Journal* 39: 461–71.
- Spies, T. A., P. A. Stine, R. A. Gravenmier, J. W. Long, and M. J. Reilly. 2018. “Synthesis of Science to Inform Land Management Within the Northwest Forest Plan area.” Gen. Tech. Rep. PNW-GTR-966. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station. 1020 p.
- Steel, Z. L., B. M. Collins, D. B. Sapsis, and S. L. Stephens. 2021. “Quantifying Pyrodiversity and its Drivers.” *Proceedings of the Royal Society B: Biological Sciences* 288: 20203202.
- Steel, Z. L., A. M. Fogg, R. Burnett, L. J. Roberts, and H. D. Safford. 2022. “When Bigger isn’t Better—Implications of Large High-Severity Wildfire Patches for Avian Diversity and Community Composition.” *Diversity and Distributions* 28(3): 439–53.
- Steel, Z. L., D. Foster, M. Coppoletta, J. M. Lydersen, S. L. Stephens, A. Paudel, S. H. Markwith, K. Merriam, and B. M. Collins. 2021. “Ecological Resilience and Vegetation Transition in the Face of Two Successive Large Wildfires.” *Journal of Ecology* 109: 3340–55.
- Steel, Z. L., G. M. Jones, B. M. Collins, R. Green, A. Koltunov, K. Purcell, S. C. Sawyer, et al. 2022a. “Code in Support of ‘Mega-Disturbances Cause Rapid Decline of Mature Conifer Forest Habitat in California’.” <https://doi.org/10.5281/zenodo.7044555>.
- Steel, Z. L., G. M. Jones, B. M. Collins, R. Green, A. Koltunov, K. L. Purcell, S. C. Sawyer, et al. 2022b. “Data in Support of ‘Mega-Disturbances Cause Rapid Decline of Mature Conifer Forest Habitat in California’.” <https://doi.org/10.6078/D1W43V>.
- Steel, Z. L., M. J. Koontz, and H. D. Safford. 2018. “The Changing Landscape of Wildfire: Burn Pattern Trends and Implications for California’s Yellow Pine and Mixed Conifer Forests.” *Landscape Ecology* 33: 1159–76.
- Steel, Z. L., H. D. Safford, and J. H. Viers. 2015. “The Fire Frequency-Severity Relationship and the Legacy of Fire Suppression in California Forests.” *Ecosphere* 6: art8.
- Stephens, S. L., A. A. Bernal, B. M. Collins, M. A. Finney, C. Lautenberger, and D. Saah. 2022. “Mass Fire Behavior Created by Extensive Tree Mortality and High Tree Density Not Predicted by Operational Fire Behavior Models in the Southern Sierra Nevada.” *Forest Ecology and Management* 518: 120258.
- Stephens, S. L., B. M. Collins, E. Biber, and P. Z. Fulé. 2016. “U.S. Federal Fire and Forest Policy: Emphasizing Resilience in Dry Forests.” *Ecosphere* 7: e01584.
- Stephens, S. L., B. M. Collins, C. J. Fettig, M. A. Finney, C. M. Hoffman, E. E. Knapp, M. P. North, H. Safford, and R. B. Wayman. 2018. “Drought, Tree Mortality, and Wildfire in Forests Adapted to Frequent Fire.” *Bioscience* 68: 77–88.
- Stephens, S. L., L. N. Kobziar, B. M. Collins, R. Davis, P. Z. Fulé, W. Gaines, J. Ganey, et al. 2019. “Is Fire “for the Birds”? How Two Rare Species Influence Fire Management across the US.” *Frontiers in Ecology and the Environment* 17: 391–9.
- Stephens, S. L., J. D. Miller, B. M. Collins, M. P. North, J. J. Keane, and S. L. Roberts. 2016. “Wildfire Impacts on California Spotted Owl Nesting Habitat in the Sierra Nevada.” *Ecosphere* 7: e01478.
- Stephens, S. L., S. Thompson, G. Boisramé, B. M. Collins, L. C. Ponisio, E. Rakhmatulina, Z. L. Steel, J. T. Stevens, J. W. van Wagtenonk, and K. Wilkin. 2021. “Fire, Water, and Biodiversity in the Sierra Nevada: A Possible Triple Win.” *Environmental Research Communications* 3: 081004.
- Stephens, S. L., A. L. Westerling, M. D. Hurteau, M. Z. Peery, C. A. Schultz, and S. Thompson. 2020. “Fire and Climate Change: Conserving Seasonally Dry Forests Is Still Possible.” *Frontiers in Ecology and the Environment* 18: 354–60.
- Stephenson, N. L., A. J. Das, N. J. Amperssee, B. M. Bulaon, and J. L. Yee. 2019. “Which Trees Die during Drought? The Key Role of Insect Host-Tree Selection.” *Journal of Ecology* 107: 2383–401.
- Stephenson, N. L., A. J. Das, R. Condit, S. E. Russo, P. J. Baker, N. G. Beckman, D. A. Coomes, et al. 2014. “Rate of Tree Carbon Accumulation Increases Continuously with Tree Size.” *Nature* 507: 90–3.
- Stevens, J. T., B. M. Collins, J. D. Miller, M. P. North, and S. L. Stephens. 2017. “Changing Spatial Patterns of Stand-Replacing Fire in California Conifer Forests.” *Forest Ecology and Management* 406: 28–36.
- Stillman, A. N., T. J. Lorenz, P. C. Fischer, R. B. Siegel, R. L. Wilkerson, M. Johnson, and M. W. Tingley. 2021. “Juvenile Survival of a Burned Forest Specialist in Response to Variation in Fire Characteristics.” *Journal of Animal Ecology*. 90: 1317–27.
- Stillman, A. N., R. B. Siegel, R. L. Wilkerson, M. Johnson, and M. W. Tingley. 2019. “Age-Dependent Habitat Relationships of a Burned Forest Specialist Emphasize the Role of Pyrodiversity in Fire Management.” *Journal of Applied Ecology* 56: 880–90.
- Stoetzel, H. J., N. P. Leseberg, S. A. Murphy, M. E. Andrew, K. J. Plant, G. N. Harrington, and J. E. M. Watson. 2020. “Modelling the Habitat of the Endangered Carpenterian Grasswren (*Amytornis dorotheae*): The Importance of Spatio-Temporal Habitat Availability in a Fire Prone Landscape.” *Global Ecology and Conservation* 24: e01341.
- Sunstein, C. R. 2002. “Beyond the Precautionary Principle Symposium: Preferences and Rational Choice: New Perspectives and Legal Implications.” *University of Pennsylvania Law Review* 151: 1003–58.
- Swetnam, T. W. 1993. “Fire History and Climate Change in Giant Sequoia Groves.” *Science* 262: 885–9.
- Taylor, A. H. 2004. “Identifying Forest Reference Conditions on Early Cut-over Lands, Lake Tahoe Basin, USA.” *Ecological Applications* 14: 1903–20.
- Tempel, D. J., R. J. Gutiérrez, J. J. Battles, D. L. Fry, Y. Su, Q. Guo, M. J. Reetz, et al. 2015. “Evaluating Short- and Long-Term Impacts of Fuels Treatments and Simulated Wildfire on an Old-Forest Species.” *Ecosphere* 6: 1–18.
- Tempel, D. J., J. J. Keane, R. J. Gutiérrez, J. D. Wolfe, G. M. Jones, A. Koltunov, C. M. Ramirez, et al. 2016. “Meta-Analysis of California Spotted Owl (*Strix Occidentalis Occidentalis*) Territory Occupancy in the Sierra Nevada: Habitat Associations and their Implications for Forest Management.” *The Condor* 118: 747–65.

- Thompson, C. M., H. Smith, R. Green, S. Wasser, and K. Purcell. 2021. "Fisher Use of Postfire Landscapes: Implications for Habitat Connectivity and Restoration." *Western North American Naturalist* 81: 225–42.
- Thompson, C. M., W. J. Zielinski, and K. L. Purcell. 2011. "Evaluating Management Risks Using Landscape Trajectory Analysis: A Case Study of California Fisher." *The Journal of Wildlife Management* 75: 1164–76.
- Tucker, J. M., M. K. Schwartz, R. L. Truex, S. M. Wisely, and F. W. Allendorf. 2014. "Sampling Affects the Detection of Genetic Subdivision and Conservation Implications for Fisher in the Sierra Nevada." *Conservation Genetics* 15: 123–36.
- USDA. 2004. "Sierra Nevada forest plan amendment. Final Supplemental Environmental Impact Statement." San Francisco, CA: USDA Forest Service, Pacific Southwest Region.
- USDA. 2019. *Conservation Strategy for the California Spotted Owl (Strix occidentalis occidentalis) in the Sierra Nevada*. San Francisco, CA: USDA Forest Service, Pacific Southwest Region.
- USDA. 2020. *2019 Aerial Survey Results: California*. Davis, CA: USDA Forest Service, Pacific Southwest Region.
- USFWS. 2020. *Endangered and Threatened Wildlife and Plants: Endangered Species Status for Southern Sierra Nevada Distinct Population Segment of Fisher*. Washington, D.C.: US DOI.
- Vanbianchi, C. M., M. A. Murphy, and K. E. Hodges. 2017. "Canada Lynx Use of Burned Areas: Conservation Implications of Changing Fire Regimes." *Ecology and Evolution* 7: 2382–94.
- van Wagtenonk, J. W., N. G. Sugihara, S. L. Stephens, A. E. Thode, K. E. Shaffer, and J. A. Fites-Kaufman. 2018. *Fire in California's Ecosystems*. Oakland, CA: University of California Press.
- Welch, K. R., H. D. Safford, and T. P. Young. 2016. "Predicting Conifer Establishment Post Wildfire in Mixed Conifer Forests of the North American Mediterranean-Climate Zone." *Ecosphere* 7: e01609.
- Westerling, A. L. 2016. "Increasing Western US Forest Wildfire Activity: Sensitivity to Changes in the Timing of Spring." *Philosophical Transactions of the Royal Society B: Biological Sciences* 371: 20150178.
- Williams, A. P., B. I. Cook, and J. E. Smerdon. 2022. "Rapid Intensification of the Emerging Southwestern North American Megadrought in 2020–2021." *Nature Climate Change* 12: 232–4.
- Williams, J., H. Safford, J. Tangenberg, Z. Steel, and A. Paulson. in revision. "High Severity Burn Area and Proportion Exceed Historic Conditions in Forests of Sierra Nevada and Adjacent Ranges (USA)."
- Wood, C. M., R. J. Gutiérrez, J. J. Keane, and M. Z. Peery. 2020. "Early Detection of Rapid Barred Owl Population Growth within the Range of the California Spotted Owl Advises the Precautionary Principle." *The Condor* 122: duz058.
- Wood, C. M., S. A. Whitmore, R. J. Gutiérrez, S. C. Sawyer, J. J. Keane, and M. Z. Peery. 2018. "Using Metapopulation Models to Assess Species Conservation–Ecosystem Restoration Trade-Offs." *Biological Conservation* 224: 248–57.
- Young, D. J. N., M. Meyer, B. Estes, S. Gross, A. Wuenschel, C. Restaino, and H. D. Safford. 2020. "Forest Recovery Following Extreme Drought in California, USA: Natural Patterns and Effects of Pre-Drought Management." *Ecological Applications* 30: e02002.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Steel, Zachary L., Gavin M. Jones, Brandon M. Collins, Rebecca Green, Alexander Koltunov, Kathryn L. Purcell, Sarah C. Sawyer, et al. 2022. "Mega-Disturbances Cause Rapid Decline of Mature Conifer Forest Habitat in California." *Ecological Applications* e2763. <https://doi.org/10.1002/eap.2763>