

CHAPTER 2

Taxonomy of Hawaiian *Cydia* (Lepidoptera: Tortricidae),
including descriptions of eight new species.

ABSTRACT

Thirteen species of endemic *Cydia* have been described from the Hawaiian Islands between 1881 and 1932 under various genera. Descriptions were based on wing colors and patterns and noted distributions and host-plant affinities where known. Hawaiian *Cydia* has a small number of wing pattern elements that are common across most species and vary within species making specific determinations problematic. The taxonomic affinities of these species were not well-understood until brought together under the genus *Cydia* and presented with genitalia dissections in the Insects of Hawaii series. Males of most Hawaiian *Cydia* possess a ventral pouch on the hindwing surface and a general reduction in features of the genitalia. Females of Hawaiian *Cydia* display greater variation in genitalia morphology, particularly in the shape of the antrum and lamellae postvaginalis. Here I describe eight new species, (*Cydia mauiensis* n.sp., *C. velocilimitata* n.sp., *C. haleakalaensis* n.sp., *C. makai* n.sp., *C. koaiae* n.sp., *C. hawaiiensis* n.sp., *C. acaciavora* n.sp., and *C. anomalosa* n.sp.), and redescribe the original thirteen based on wing markings, presents or absence of the hindwing pouch, and features of both male and female genitalia. Larvae of Hawaiian *Cydia* feed on reproductive structures such as seeds, flowers, or buds and within twigs, under bark, or in rust galls of endemic Hawaiian plants in the family Fabaceae. Distributions, natural enemies, and host-plant associations are presented where known.

INTRODUCTION

Cydia Hübner, a worldwide genus in the tribe Grapholitini (Olethreutinae), is represented by at least 21 species in the Hawaiian Islands. Found from shoreline to treeline on the main Hawaiian Islands (Kauai, Oahu, Molokai, Maui, and Hawaii), *Cydia* plays an important role in native forest ecosystems. The larvae of Hawaiian *Cydia* are associated with the native plants *Acacia koa* A.Gray, *A. koaia* Hillebr., *Canavalia* spp., *Sophora chrysophylla* (Salis.) Seem., *Strongylodon ruber* Vogel, and *Vicia menziesii* Spreng. in the family Fabaceae, feeding in generative structures (Figures 1-8) such as seeds, flowers, terminal buds, and under bark (Swezey 1954, Zimmerman 1978). As seed predators, Hawaiian *Cydia* can have a dramatic impact on the contributions of particular plant species to the seed bank (Zimmerman 1978). By feeding under the bark of dead and dying branches, larvae expose the wood to other invertebrate colonizers and decay fungi. And for several rare and endangered forest bird species in Hawaii that glean insects from seeds, flowers, and under bark, *Cydia* are an important protein-rich food resource (Banko et al. 2002, Pratt et al. 2009).

The appreciation of *Cydia* species diversity in Hawaii has been hampered, however, by a lack of conspicuous taxonomically useful characters and by extreme polymorphism in wing color patterns (Figures 9-12). The genus *Cydia* is characterized by a reduction in ornamentation in the male genitalia (i.e. loss of uncus, socii, and gnathos); characters that are often diagnostic for species in other groups of Lepidoptera (Komai 1999). Although subtle differences in the shape of the male valvae, aedeagus, and tegumen, and the female antrum and “ostial plate” (lamellae ante- and postvaginalis) can be used for identification, careful preparation and examination of genitalia are necessary to appreciate these differences. Conversely, whereas many Lepidoptera species have very characteristic wing patterns, some Hawaiian *Cydia* species vary widely in both color and wing pattern elements. The three most variable species, *C. plicata*, *C. makai*, and *C.*

walsinghamii, also have the broadest distribution in the islands, making superficial identifications problematic.

The thirteen named endemic *Cydia* species recognized from Hawaii were described between 1881 and 1932, and appear to be closely allied (Zimmerman 1978). Butler (1881, 1882) described the first Hawaiian species, *C. rufipennis* and *C. walsinghamii*, from the island of Oahu based on collections by the reverend Thomas Blackburn. Walsingham (1907) later described *C. conspicua*, *C. crassicornis*, *C. falsifalcella*, *C. latifemoris*, *C. montana*, *C. obliqua*, *C. plicata* and *C. storeella* from the monumental Fauna Hawaiiensis collections of R.C.L. Perkins (1913). Meyrick (1932), in his Exotic Microlepidoptera series, described three more species, *C. chlorostola*, *C. gypsograptus*, and *C. parapteryx*. An additional species, *Cydia* “new species 1”, was known from collections before the publication of the Insects of Hawaii microlepidoptera volume, but remained undescribed (Zimmerman 1978). Several species are known from only one to three individuals used in the original descriptions: *C. chlorostola* (1♀), *C. crassicornis* (2♂), *C. gypsograptus* (1♂), *C. obliqua* (3♀), and *C. storeella* (1♀). Although each of these appears to be unique species, the lack of specimens is a concern because of known variation in wing color patterns, which were the basis of all the original descriptions.

The affinity of the Hawaiian species to *Cydia* outside of Hawaii remains obscured (Oboyski Chapter 3). Butler (1881) described *C. rufipennis* in the genus *Phoxopterus* Treitschke (= *Ancylis* Hübner), which is assigned to the tribe Enarmoniini. A year later, Butler (1882) described *C. walsinghamii* in the genus *Proteopteryx* Walsingham (= *Epinotia* Hübner), which is in the tribe Eucosmini. Walsingham (1907) proposed a new genus, *Adenoneura*, based on the presence of a glandular pouch along the cubital vein in the hindwing of males (*Adeno* = gland, *neura* = nerve) (Figures 13-18). He placed four of his new species in this genus, stating it was allied to *Thiodia* Hübner (Eucosmini), and tentatively placed the rest, (males lacking the glandular pouch or species known only from females), in *Enarmonia* Hübner (Enarmoniini). Meyrick (1932) was the first to recognize the affinity of Hawaiian species to other *Cydia* when he described *C. chlorostola* in the genus *Laspeyresia* Hübner (= *Cydia*). The use of genitalic characters in microlepidopterology did not come into vogue until the early 20th century. All previous descriptions had been based on external features, particularly wing venation and color patterns. Zimmerman (1978) presented genitalia preparations of the type specimens of Hawaiian *Cydia* and brought the twelve (plus one undescribed) species together under the name *Cydia* and tentatively suggested they derived from a broadly distributed legume-feeder.

Here I describe eight new species of Hawaiian *Cydia* and redescribe the thirteen previously known species. Previous treatments of Hawaiian *Cydia* are inconsistent in their descriptions and none includes descriptions of genitalia. A comprehensive treatment of the currently known species will 1) provide a common framework for identifying specimens and recognizing new species, and 2) synthesize our current knowledge of Hawaiian *Cydia* diversity.

MATERIALS AND METHODS

Material Examined

I examined freshly prepared field-caught specimens as well as historical material from several museum collections, as follows:

BPBM	Bernice P. Bishop Museum, Honolulu, Hawaii
BMNH	The Natural History Museum, London
CAS	California Academy of Sciences, San Francisco, California
EMEC	Essig Museum of Entomology, University of California, Berkeley
HAVO	Hawaii Volcanoes National Park, Volcano, Hawaii
HDOA	Hawaii Department of Agriculture, Honolulu, Hawaii
MEM	Mississippi Entomology Museum
UHMA	University of Hawaii at Manoa, Honolulu, Hawaii
USNM	United States National Museum, Washington, D.C.

Holotypes of all previously described Hawaiian *Cydia* species and slide preparations of their genitalia are housed at the Natural History Museum, London. Well-prepared series of Hawaiian *Cydia* collected by Dr. Klaus Sattler in 1973, 1976, and 1982 are not yet integrated into the BMNH collections but were available for examination. The Bernice P. Bishop Museum and Hawaii Department of Agriculture collections each hold paratypes and short series of some species, and the Hawaii Volcanoes National Park and University of Hawaii, Manoa collections each hold short series of particular species. A small number of specimens exist in other institutional and private collections, such as the California Academy of Sciences, the Mississippi Entomology Museum, and the United States National Museum. The remainder of the material examined I collected and prepared between 1998 and 2008. Voucher specimens were deposited in the BPBM, HDOA, and EMEC.

Field collections

Hawaiian *Cydia* species were collected on all the main islands (Hawaii, Maui, Molokai, Oahu, and Kauai), but not from Kahoolawe, Lanai, and Niihau. Because the Island of Kahoolawe was virtually denuded of native vegetation by ungulate grazing, it is unlikely that any native *Cydia* survive there. Five to ten *Sophora chrysophylla* trees remain on the island of Lanai along with planted *Acacia koa* trees, but I failed to find larval or moth specimens. However, examination of historic collections from Lanai of *S. chrysophylla* in the Bishop Museum herbarium revealed emergence holes from seedpods similar to *Cydia* emergence holes from other islands. I also visited the island of Nihoa, the youngest and most pristine in the Northwest Hawaiian Islands chain, but found no *Cydia* despite the presence of an abundant potential host legume, *Sesbania tomentosa* Hook. & Arn.

Field-collected specimens were taken either as adults at ultraviolet lights or as larvae from host plant material. Adults collected at lights were killed either in the field with potassium cyanide, or in a refrigerator freezer (-20°C). Prior to pinning and spreading adults, I removed and placed into 95% EtOH 1-3 legs from a subset of moths for subsequent DNA analysis (Oboyski Chapter 3). Field collected larvae were reared in plastic containers fitted with mesh lids (Brenner et al. 2002) in order to confirm host-plant and parasitoid relationships.

Dissections

Zimmerman (1978) dissected the genitalia of type specimens housed at the BMNH. I examined each of these preparations and found certain features difficult to discern due to the orientation and weak staining of the genitalia. Zimmerman (1978) states, “there are obvious differences in the genitalia, but they are not clearly shown in all of my illustrations.” Therefore, where possible, I made additional dissections from conspecific specimens. I prepared dissections following Robinson (1976) as follows: Abdomens were removed from the thorax of dry specimens and briefly “wetted” in 70% EtOH and placed in sub-boiling 10% KOH for 3 to 5 minutes, or until sclerotized tissues were pliable. The entire abdomen was then placed in distilled water for cleaning which consisted of brushing away scales with an ultra-fine paintbrush, gently “squashing” the abdomen with blunt soft forceps to squeeze out macerated soft tissues, and flushing with distilled water using a fine syringe. Male genitalia were pulled free from the rest of the abdomen (or “pelt”) with forceps, naturally separating between the 8th and 9th segments. As the female lamellae antevaginalis and postvaginalis (or “ostial plate”) are located on the 8th segment, female genitalia were pulled free from the pelt between the 7th and 8th segments. Further cleaning of loose tissue and scales was done in distilled water using a fine brush and curled tip of a minuten pin. Both the pelt and genitalia were then dipped for one to two minutes in concentrated Chlorozol Black E stain and rinsed in distilled water. Next the pelt and genitalia were dipped in concentrated Eosin-Y stain for approximately 5-10 minutes then rinsed with distilled water. The combination of Chlorozol Black E and Eosin-Y creates a contrast between the weakly sclerotized tissues (stained blue-gray by the Chlorozol Black E) and the more-strongly sclerotized tissues (stained red by the Eosin-Y). Next the genitalia were positioned in a series of baths of 70%, then 95% EtOH, for 10-15 minutes, often held in position by a small shard of microscope slide. Genitalia were then placed in Euparal Essence[®] until mounted in Euparal[®] mounting medium (BioQuip Products, Rancho Dominguez, CA) on glass slides with cover slips.

The juxta of male *Cydia* genitalia, like those of many Olethreutinae, is fused with the caulis and anellus forming a stout, inflexible substrate to which the aedeagus is attached (Horak 2006). This poses a challenge in positioning the valvae and aedeagus, resulting in features, particularly the valvae, aedeagus, and tegumen, being distorted or in different planes of focus. For this reason it was sometimes preferable to separate the valvae and aedeagus before mounting them on glass slides. It is also common practice in the study of some lepidopteran taxa to evert the vesica of the aedeagus in order to better view its shape and any cornuti present. None of the specimens examined appeared to have cornuti (deciduous or otherwise), so no attempt was made to evert the vesica to avoid potentially damaging the preparations.

Wings were prepared for examination of venation patterns following Zimmerman (1978). Wings (usually right side) were broken or cut free from the body and “wetted” with 70% EtOH. The wings were then placed in pure household bleach (6% sodium hypochlorite) for 30-60 seconds, or until small bubbles began to form. The now soft and delicate wings were transferred to distilled water for cleaning with a fine brush and the hooked tip of a minuten pin. Wings were then stained with acid fuchsin to visualize the veins. The stained wings were then rinsed in 70% EtOH and passed through 95% EtOH to drive away excess moisture. Still in 95% EtOH, the

wings were then floated onto glass slides and mounted with Euparal[®], once the EtOH had evaporated to the edges of the wings.

Illustrations

Slide preparations (wings and genitalia) were photographed with a Leica[®] digital camera and microscope using Automontage[®] software (Syncroscopy, Cambridge, UK and Maryland, USA). These digital images were then traced and shaded in Adobe Illustrator[®] and Photoshop[®] (San Jose, California, USA). Similarly, figures from Zimmerman (1978) were redrawn after examining the original slides and taking note of features difficult to discern in the published photographs.

Morphological Characters

The following characters were considered for diagnosing and describing *Cydia* species.

Colors: Descriptive colors follow the language of the original descriptions for consistency and comparison, and can be translated as follows: buff (yellowish beige), cinereous (ash-gray), ferruginous (reddish-brown, rust), fuliginous (sooty, dusky), fuscous (dark gray, grayish-brown), ochreous (earthy reddish-yellow), olivaceous (olive green, dusky yellowish green), plumbaginous (lead gray), testaceous (dull brick red).

Scales: Many species have head, thorax, and leg scales with light-colored tips that contrast with their darker bases. Although not used here as a diagnostic character, the presence of these contrasting scales contributes to the overall color patterns.

Head: The antennae, palpi, and vestiture of the head are typical of most *Cydia* species and consistent for all Hawaiian species, and therefore not of diagnostic value.

Thorax: Wing expanse (exp.) was measured to the nearest half-millimeter from wing tip to wing tip (including fringe) of fully spread specimens. Wing expanse is variable and often related to larval food quantity and quality (Miller 1990) but is diagnostic for some species. Wing venation is nearly uniform for all Hawaiian *Cydia* species (Figure 9), except as noted where the presence of a hindwing pouch displaces vein A3 towards the anal area. Scale color of the thorax and banding patterns on the legs are often useful in confirming identifications, but not consistent enough to be diagnostic. Wing patterns are limited to a small number of pattern elements (Figure 10) that are variably present or absent on individuals within a species. The combination of pattern elements and their colors can be used to identify many species, but with caution because of the high degree of polymorphism found within many species. Species within the *Canavalia*-feeding group all appear to have a sinuous termen and crescent-shaped apical patch on the forewing (Figure 11) as opposed to termen straight and subtriangular-shaped apical patch of other Hawaiian *Cydia* (Figure 12). The absence of a ventral pouch within the hindwing of males (Figure 13) is diagnostic for four species, and the presence of an anal roll in male hindwings is known only in *C. anomalosa* n.sp. (Figures 19-24). Variations in wing patterns are illustrated by photographs (Figures 25-51).

Abdomen: Male (Figures 52-69) and female (Figures 70-86) genitalia have the most consistent diagnostic value in Hawaiian *Cydia*. The invagination of the ventral margin at the base of the male cucullus is variable among species but of limited value, and potentially deceptive, because perceiving it is dependent on the orientation of the genitalia in the slide preparation. Similarly, the shape of the caudal ridge, or crista, of the tegumen (in the absence of a developed uncus) can be diagnostic for some species when the preparation allows the proper view of this structure. For males, the most diagnostic feature is the tip of the aedeagus which may be flared, spatulate, or

excavated. For females, the shape of the lamellae postvaginalis and antrum (i.e. basal sclerotization at the junction of the ductus bursae with the ostium) are diagnostically more informative than characters in the male genitalia for this group. Zimmerman (1978) noted that the proximity of the signa in the female corpus bursae was a character indicative of some species, but closer inspection of his slide preparations revealed this was an illusion caused by orientation of signa on opposite sides of the corpus bursae.

Genus *Cydia* Hübner, 1825

Cydia species are easily separated from other Hawaiian Tortricidae using the keys in Zimmerman (1978) based on wing venation, labial palpi, and some secondary sexual characters. The synonymies and general morphology of the genus *Cydia* are reviewed elsewhere (Oboyski Chapter 1, Komai 1999, Komai and Horak 2006). Hawaiian *Cydia* conforms to the general description of other *Cydia* with a few notable exceptions. Males of Hawaiian *Cydia* lack the hindwing anal roll and associated sex scales (Figures 22, 24), and thecae along hindwing vein A3 (Figure 20) found in most other *Cydia* species (Komai and Horak 2006), with the exception of *C. anomalousa* n.sp., which possess an anal roll and phylliform scales only (Figures 21, 23). Where males are known, 15 of 18 species of Hawaiian *Cydia* have a glandular ventral pouch that opens dorsally along the path of the CuP vein (Figures 9, 13, 14), which encloses modified cubital pecten scales (Figure 15, 17). A similar structure occurs in *C. latiferreana* (Walsingham) (Figures 16, 18), *C. maackiana* (Danilevsky), and several other *Cydia* species to a lesser degree, but appear to be derived independently in Hawaiian *Cydia* based on the microstructure of the modified pecten scales (Figures 15-18). The presence of the hindwing ventral pouch is accompanied by a shift in vein A3 towards the anal margin (Figure 9).

Cydia chlorostola (Meyrick 1932) (Figure 70)

Laspeyresia chlorostola Meyrick 1932:226

Cydia chlorostola. – Zimmerman 1978:585, Figures 378 (moth), 389 (genitalia)

DIAGNOSIS: The most pale of all the Hawaiian *Cydia*, nearly uniform whitish. Distinguished from other Hawaiian *Cydia* by the complexity of the antrum sclerotization (Figure 70). Most similar to, and likely to be confused with, very pale forms of the *Canavalia*-feeding Hawaiian species (*C. falsifalcella*, *C. mauiensis*, *C. parapteryx*, *C. velocilimitata*), which have similarly complex antra (Figures 71-73).

DESCRIPTION: (exp. 16 mm, n=1) **Head:** Antennae and labial palpi uniformly buff-white. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Head buff-white, slightly darker laterally. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally and ventrally buff-white (holotype partly denuded), tegulae somewhat lighter. Legs uniformly white-cinereous. No discernable sex scales (e.g. hair pencils). **Forewings:** Slightly dilated distally, costa gently arched, apex obtuse, termen somewhat sinuate. Nearly uniform buff-white with no distinguishable striae or fascia, ocellus consisting of three black dots of one scale each, bordered distally by faint silvery-whitish line, fringe white-cinereous. Ventrally somewhat

darker ochreous-white basally grading to buff-white distally. **Hindwings:** White-cinereous, faintly gray along fold and above cell. Fringe and ventral hindwing uniformly white-cinereous. **Abdomen:** (removed for dissection). **Male genitalia:** (male not known). **Female genitalia:** (Figure 70) Lamellae postvaginalis nearly as broad as long and somewhat centrally constricted, antrum with complex sclerotization pattern, extending well beyond the junction with ductus bursae. Corpus bursae with diverticulum, two long falcate signa with rounded tips. Ductus bursae slightly longer than width of corpus bursae, tapering gradually from corpus bursae to ostium.

TYPE MATERIAL: *Laspeyresia chlorostola* Meyrick – ♀ holotype (BMNH): Hawaiian Islands, Oahu, Waialua, P. 09 [RCL Perkins 1909]; genitalia slide BM 9546 Clarke.

ADDITIONAL MATERIAL: The female type is the only known specimen of this species.

BIOLOGY: There is no information available regarding larval biology, host plants, habitat, predators, or parasitoids.

DISTRIBUTION: Perkins' label includes only the general description of "Waialua, Oahu." Meyrick (1932) suggested this species was introduced along with a legume host, probably of Asiatic origins, but is not known from any other locations. Zimmerman (1978) disagreed and considered the species endemic to Hawaii. In the absence of more specimens I decline to conjecture any further than to suggest this is an endemic Hawaiian species.

REMARKS: This species is probably extinct. The type location, Waialua region of Oahu, no longer contains native habitats, and sampling of the nearby native forests has not produced any new specimens. Aspects of the female genitalia (see description above) suggest that this is a member of the Hawaiian *Cydia* group, and not an introduced species as Meyrick suggests.

Cydia gypsograpt (Meyrick 1932)
(Figure 52)

Adenoneura gypsograpt Meyrick 1932:222

Cydia gypsograpt. – Zimmerman 1978:591; Figures 379 (moth), 384 (♂ genitalia)

DIAGNOSIS: A mostly white species with dark gray strigulae and pretornal blotch, but without costal triangle and distinct medial fascia common in other Hawaiian species. Male genitalia difficult to discern from *Canavalia*-feeding species group (*C. falsifalcella*, *C. mauiensis*, *C. parapteryx*, *C. velocilimitata*).

DESCRIPTION: (exp. 13 mm, n=1) Colored scales on head, body, and legs mostly shades of grayish-brown with whitish tips. **Head:** Antennae brownish-cinereous, head and labial palpi dorsum white, palpi with fuscous lateral streak. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally and ventrally generally whitish, mid-dorsum light brown scales with white tips, tegulae light brown anteriorly grading to longer white scales posteriorly. Without dorsal tuft of scales. Legs whitish, foretibia banded brown and white, midtibia less so. No discernable sex scales (e.g. hair pencils). **Forewings:** Slightly dilated distally, costa gently arched, apex obtuse, termen

somewhat sinuate. Ground color buff white. Costal strigulae distinct, directed distally towards termen, striae fragmented through cell but prominent again from fold to dorsal margin as dark streaks between dorsal strigulae. No triangular costal patch. Distinct, broadly rounded-triangular fuscous pretornal blotch. Pale ocellar area with one or two distinct ocellar spots, bounded distally by light fuscous then silvery crescent. Apex with a fuscous patch probably extending into fringe (fringe worn away). Continuation of fuscous stria to central termen extends into fringe. Ventrally uniform light fuscous. **Hindwings:** Fuscous, lighter anteriorly, with silvery-white fringe. Ventrally uniform light fuscous. Males with glandular ventral pouch below cubital vein opening dorsally and enclosing elongate modified pecten scales. Vein A3 in male displaced towards the anal margin. Males without an anal role or androconial scales (thecae) along vein A3. **Abdomen:** (abdomen removed). **Male genitalia:** (Figure 52) Tegumen simple with broadly rounded caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa gently concave, ventral margin with moderate invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with dense short setae along ventral and distal margin, interspersed with fewer long setae. Aedeagus simple, curved, without cornuti, spatulate tip excavated dorsally for 1/5th of length of aedeagus. **Female genitalia:** (females not known).

TYPE MATERIAL: *Adenoneura gypsograpt* – ♂ holotype (BMNH): Oahu, Honolulu, P. 08 [RCL Perkins 1908]; genitalia slide BM 9543 Clarke.

ADDITIONAL MATERIAL: This species is known only from the male holotype.

BIOLOGY: There is no information available regarding larval biology, host plants, habitat, predators, or parasitoids.

DISTRIBUTION: Endemic to Hawaiian Islands: This species is known only from a single specimen collected by RCL Perkins near Honolulu, Oahu.

REMARKS: This species is probably extinct. The Honolulu region of Oahu no longer contains native habitats, and sampling of nearby native forests has not turned up any new specimens. The presence of the hindwing pouch in the male suggests that this is a member of the Hawaiian *Cydia* group.

Cydia mauiensis n. sp.
(Figures 11, 25, 53, 71)

DIAGNOSIS: Most closely allied with *C. falsifalcella*, *C. parapteryx*, and *C. velocilimitata*, being the Maui Island taxon of the *Canavalia*-feeding species group, with wing patterns similar. Differentiated from these other species by its distribution and the shape of the female antrum (Figure 71).

DESCRIPTION: (exp. 15-17 mm, n=5) **Head:** Antennae, head, and labial palpi light ochreous. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally light ochreous, tegulae same. Without dorsal tuft of scales. Ventrally lighter. Legs light brown-ochreous. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figures 11, 25) Slightly dilated distally, costa gently arched, apex

obtuse, termen somewhat sinuous. Ground color light ochreous. Costal strigulae distinct, silvery light brown basally, whitish-buff distally, directed towards termen, striae indistinct except near apex. Basal area light brown-ochreous to buff, mottled with occasional brown or fuscous scales. No triangular costal patch. Oblique medial fascia extending from end of cell towards basal dorsum brown-ochreous with darker brown distal border. Whitish discal patch present and extending along medial fascia. Brown-ochreous pretornal patch vaguely present. Ocellar patch light ochreous with three distinct ocellar spots, bordered distally by a silvery-white crescent. Apex with wide crescent-shaped light brown-ochreous patch extending into fringe as dark brown-ochreous. Continuation of light brown-ochreous stria to central termen extends as dark brown-ochreous into fringe. Fringe otherwise light brown-ochreous. Ventrally uniform brown-cinereous. **Hindwings:** Dorsal and ventral light bronze-ochreous, somewhat lighter basally and ventrally. Males with glandular ventral pouch along path of CuP, opening dorsally and enclosing elongate modified pecten scales. Vein A3 in males displaced towards anal margin. Males without anal roll or androconial scales (thecae) along vein A3. **Abdomen:** light brown-ochreous dorsally, lighter ventrally. **Male genitalia:** (Figure 53) Tegumen simple with bilobed crista along caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa gently concave, ventral edge with somewhat deep invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with dense short setae along ventral and distal margin interspersed with longer setae distally. Aedeagus simple, curved, without cornuti, and with tip spatulate and excavated dorsally for approximately one-third the length of aedeagus. **Female genitalia:** (Figure 71) Lamellae postvaginalis nearly as wide as long and slightly constricted centrally. Antrum with prominent dorsal lobe near ostium and ventral lobe extending well beyond the dorsal junction with ductus bursae. Corpus bursae with diverticulum, two moderately long falcate signa. Ductus bursae shorter than width of corpus bursae, slightly wider at junction with corpus bursae than junction with antrum.

TYPE MATERIAL: *Cydia mauiensis* - ♂ holotype (BPBM): Hawaiian Islands, Maui, Kipahulu Valley, Oheo campground, 10.VII.2005, PT Oboyski; UV light; Oboyski specimen PTO-544.1. ♀ allotype (BPBM): Hawaiian Islands, Maui, Kipahulu Valley, Oheo campground, 10.VII.2005, PT Oboyski; UV light; Oboyski specimen PTO-544.4; genitalia slide PTO-s154. 7♂ paratypes (EMEC): Hawaiian Islands, Maui, Kipahulu Valley, Oheo campground, 2 m., UV light nr. *Canavalia* vines 10.VII.2005, PT Oboyski. Oboyski specimen PTO-544.2, genitalia slide PTO-s254.

ADDITIONAL MATERIAL: This species is known only from a single collection of nine specimens from the type locality.

BIOLOGY: Adults were collected at UV lights in proximity to *Canavalia* vines. Although no larvae were found within seedpods, like its close relatives, *C. falsifalcella*, *C. parapteryx*, and *C. velocilimitata*, this species probably feeds opportunistically within flower peduncles and stems of *Canavalia* spp.

DISTRIBUTION: Endemic to Hawaiian Islands: Maui. Possibly widespread and locally common where *Canavalia* is abundant, but known only from the type locality at Oheo campground, Haleakala National Park, Maui.

REMARKS: In molecular phylogenetic analyses (Oboyski Chapter 3), this species appears most basal in the Hawaiian *Cydia* clade or forms a basal clade with the three other *Canavalia*-feeding species.

Cydia falsifalcella (Walsingham 1907)
(Figures 26, 27, 54, 72)

Adenoneura falsifalcellum Walsingham 1907:677; Plate 10, Figure 17 (moth)

Cydia falsifalcella. – Zimmerman 1978:586; Figures 371 (head, wing venation), 373 (wing venation), 379 (moth), 384 (♂ genitalia)

DIAGNOSIS: Most closely allied with *C. mauiensis*, *C. parapteryx*, and *C. velocilimitata*, being the Hawaii Island taxon of the *Canavalia*-feeding species group, with wing patterns similar. Differentiated from these other species by its distribution and the shape of the female antrum (Figure 72).

DESCRIPTION: (exp. 13-18 mm, n=6) Colored scales on head, body, and legs mostly shades of grayish-brown with whitish tips. **Head:** Antennae and labial palpi uniformly light brown-cinereous. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Head uniformly light brown-cinereous. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally and ventrally light brown-cinereous, tegulae same. Without dorsal tuft of scales. Legs light brown-cinereous, fore- and mid-tibiae banded with fuscous bases and lighter distally, hindleg tarsi likewise banded. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figures 26-27) Slightly dilated distally, costa gently arched, apex obtuse, termen somewhat sinuate. Ground color ochreous-white, overlaid by variable pattern elements. Costal strigulae distinct, directed distally towards termen, striae vague below costa in basal area, more distinct and ferruginous distally. Occasionally with a distinct triangular costal patch suffused from base to 2/3 length of costa, posteriorly overlapping the fold, ranging in color from dark fuscous to ferruginous. Costal patch often not fully suffused anterior and basal leaving an oblique medial fascia extending from end of cell towards basal dorsum (as in ♂ holotype), bordered distally by a distinct whitish discal patch at end of cell. Light brown pretornal patch sometimes vaguely evident. Pale ocellar patch with three distinct ocellar spots, bordered distally by a silvery-white crescent. Apex with a crescent-shaped ferruginous patch extending into fringe. Continuation of ferruginous stria to central termen extends into fringe. Ventrally strigulae apparent along costal margin, otherwise uniformly brown-cinereous. **Hindwings:** Uniformly light brown-cinereous dorsally, ventrally vague light brown maculations on fuscous background. Males with glandular ventral pouch along path of CuP, opening dorsally and enclosing elongate modified pecten scales. Vein A3 in male displaced towards the anal margin. Males without an anal role or androconial scales (thecae) along vein A3. **Abdomen:** Uniformly light brown-cinereous. **Male genitalia:** (Figure 54) Tegumen simple, occasionally with single lobed crista along the caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa gently concave, ventral edge with deep invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with dense short setae along ventral and distal margin, interspersed with fewer long setae. Aedeagus simple, curved, without cornuti, flared spatulate tip excavated dorsally for one-fifth length of aedeagus. **Female genitalia:** (Figure 72) Lamellae postvaginallis slightly longer than wide, hourglass-shaped, antrum with complex sclerotization

pattern, extending beyond junction with ductus bursae. Corpus bursae with diverticulum, two long falcate signa. Ductus bursae approximately as long as width of corpus bursae, tapering gradually from corpus bursae to antrum.

TYPE MATERIAL: *Adenoneura falsifalcellum* Walsingham – ♂ holotype (BMNH): Hawaii: Olaa, VII. 1895, RCL Perkins; Walsingham specimen 27506; genitalia slide BM 2051. 1♂ paratype (BMNH): Hawaii, Olaa, ix.1896, RCL Perkins; Walsingham specimen 28043. 3♂ paratype. (BMNH): Hawaii, Olaa, 2000 ft, RCL Perkins; Walsingham specimens 28630, 28638, 28640. 2♂ (BPBM): Hawaii, Olaa, 2000 ft, xi.1896, RCL Perkins; Walsingham specimens 28642, 28648; genitalia & wings (28648), “slide 79, ecz Oct 1940.”

ADDITIONAL MATERIAL: 1♂ (BPBM): Hawaii, Mauna Loa, Kaeuhou road, 5900ft, reared: *Vicia menziesii*, 5-27.iii.1979, C Hodges; USFS #79-1; genitalia slide PTO-s182. 1♀ (BPBM): Hawaii, Mauna Loa, Keauhou road, 5900ft, reared: *Vicia menziesii*, 5-27.iii.1979, C Hodges; USFS #79-1; genitalia slide PTO-s183. 2♂ (EMEC): Hawaii, Mauna Loa, Manuka State Park, loop trail, 560 m, UV light nr. *Canavalia hawaiiensis*, 28.viii.2005, PT Oboyski; genitalia slide PTO-s251.

BIOLOGY: According to Zimmerman (1978), descriptions of larval biology attributed to *C. falsifalcella* prior to 1932, when *C. parapteryx* was described from Oahu, belong to the latter species. Probably, like *C. parapteryx*, *C. falsifalcella* larvae feed on *Canavalia* species. Inspection of the few seedpods of *Canavalia hawaiiensis* growing at Manuka, Hawaii Island showed no indication of larval feeding; however, two adults were collected at an ultraviolet light positioned amid the *Canavalia* vines. Larvae probably develop opportunistically in the flower peduncles, seedpods, or vines of this plant. While *Canavalia hawaiiensis* is rare in nature, *Canavalia galeata*, the type host of *C. parapteryx* on Oahu, is common at low elevations. However, UV light trapping and seedpod inspection of *Canavalia galeata* have not produced any *Cydia* specimens on Hawaii Island. Two specimens of *C. falsifalcella* also were reared from the seeds of *Vicia menziessii* at Keauhou, Hawaii, by C. Hodges, suggesting that this species may have a somewhat varied diet. *Vicia menziesii* is a critically endangered plant represented by less than a dozen mature, naturally growing plants, and dozens of struggling out-planted seedlings. None of these plants has set seed in many years (Kealii Bio, USGS-BRD, personal communication). However, *Vicia* plants were once more common in this area and may have been an alternate host-plant for this species, and the source for Perkins' original collections in Olaa.

DISTRIBUTION: Endemic to Hawaiian Islands: Hawaii – uncommon and very localized.

REMARKS: *C. falsifalcella* closely resembles its sister species *C. parapteryx* from Oahu and has been confused with this latter species in the literature (Zimmerman, 1978) (see notes under Biology above).

Cydia parapteryx (Meyrick 1932)
(Figures 9, 28, 55, 73)

Enarmonia sp. Swezey 1908:15

Adenoneura falsifalcella (sensu Perkins 1913:clxviii, not Walsingham 1907)

Adenoneura parapteryx Meyrick, 1932:222

Cydia parapteryx. – Zimmerman 1978:608; Figures 374 (wing venation), 381 (moth), 386 (♂ genitalia), 395 (♀ genitalia)

DIAGNOSIS: Most closely allied with *C. falsifalcella*, *C. mauiensis*, and *C. velocilimitata*, being the Oahu Island taxon of the *Canavalia*-feeding species group, with wing patterns similar. Differentiated from these other species by its distribution, the lack of excavation of the tip of the male aedeagus (Figure 55), and the shape of the female antrum (Figure 73).

DESCRIPTION: (exp. 14-24 mm, n=23) **Head:** Antennae, head, and labial palpi light brown-ochreous. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally light brown-ochreous, tegulae same. Without dorsal tuft of scales. Ventrally lighter, somewhat darker around coxae. Legs light brown-ochreous. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figure 28) Slightly dilated distally, costa gently arched, apex obtuse, termen somewhat sinuous, nearly vertical. Ground color light brown-ochreous. Costal strigulae distinct, silvery light brown basally, whitish-buff distally, directed distally towards termen, striae indistinct in basal half creating mottled appearance or obscured by suffusion of ferruginous to fuscous triangular costal patch. Oblique medial fascia along distal edge of costal triangle extending from end of cell towards basal dorsum dark ferruginous often with dark brown-fuscous distal border. Whitish discal patch present and often distinct. Pretornal patch absent or very vague. Ocellar patch mottled light and dark brown with three distinct ocellar spots, bordered distally by a silvery-white crescent. Apex with wide crescent-shaped ferruginous patch extending into fringe as fuscous-ferruginous. Continuation of ferruginous stria to central termen extends as fuscous-ferruginous into fringe. Fringe otherwise light brown-buff. Ventrally strigulae apparent along costal margin, otherwise uniform brown-cinereous. **Hindwings:** Dorsal and ventral ferruginous, somewhat lighter basally and ventrally, maculation somewhat apparent ventrally. Males with glandular ventral pouch along path of CuP, opening dorsally and enclosing elongate modified pecten scales. Vein A3 in males displaced towards anal margin. Males without anal roll or androconial scales (thecae) along vein A3. **Abdomen:** light brown-ferruginous dorsally, lighter ventrally. **Male genitalia:** (Figure 55) Tegumen simple with broadly flattened caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa gently concave, ventral edge with somewhat deep invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with dense short setae along ventral and distal margin interspersed with longer setae distally. Aedeagus simple, curved, without cornuti, and tip bilobed. **Female genitalia:** (Figure 73) Lamellae postvaginalis somewhat longer than wide, slightly constricted centrally. Antrum relatively shortened with prominent dorsal lobe near ostium and ventral lobe extending just beyond dorsal junction with ductus bursae. Corpus bursae with diverticulum, two long falcate signa. Ductus bursae shorter than width of corpus bursae, only slightly wider at junction with corpus bursae than junction with antrum.

TYPE MATERIAL: *Adenoneura parapteryx* – ♂ lectotype (BMNH): Oahu, Honolulu, P. 09 [1909], bred from *Canavalia galeata*, RCL Perkins; genitalia slide BM 9544. 2♀ paralectotype (BMNH): Oahu, Honolulu, P. 09, bred from *Canavalia galeata*, RCL Perkins; genitalia slide BM 7549. 3♂ paralectotype (BMNH): Oahu, Honolulu, P. 09 [RCL Perkins 1909], bred from *Canavalia galeata*; wing slide BM 7532.

ADDITIONAL MATERIAL: 1♀ (BPBM): Oahu, Mt Kaala, ex: *Canavalia galeata* seeds, 14.x.1928, OH Swezey. 3 (HDOA): Oahu, Mt Kaala, ex: *Canavalia galeata* seeds, (11.xi.1926, 14.x.1928, 23.i.1929), OH Swezey. 4 (HDOA): Oahu, Honolulu, (10.viii.1907, 27.i.1908, 1.ii.1908, 28.iv.1908). 1 larva (HDOA): Oahu, Honolulu, ex: *Canavalia tursida*, 31.vii.1907. 2 (HDOA): Oahu, Honolulu, ex: *Canavalia microcarpa*, 9.viii.1907. 1 (HDOA): Oahu, HPSA station, ex: beans, 22.vii.1907. 1 (HDOA): Oahu, Piko trail, ex: *Canavalia*, 30.xii.1933, OH Swezey. 1 (HDOA): Oahu, Kahaluu, light trap, vii.1969, W Au. 1 (HDOA): Oahu, Nuuanu, ex: *Strogylodon*, 1.i.1916. 3♂ (EMEC) Oahu, S. Waianae Mtns., Honouliuli TNC Preserve, Kaluaa section, 585 m., UV light nr. *Strongyloдон* vines, 24.vii.2005, PT Oboyski, Oboyski specimens PTO-563.1,2,3. 2♀ (CAS) Oahu, Mt Kaala, ex: *Canavalia galeata* seeds, 14.x.1928, OH Swezey. 2 (CAS) Oahu, Honolulu, 27.i.1908. 4 (USNM) Oahu, Honolulu.

BIOLOGY: Larvae feed in the unripe seeds, fleshy walls of developing seedpods, flower cluster peduncles, leaf petioles, and vining branches of *Canavalia galeata* (Fabaceae) (Swezey, 1908 as *Enarmonia* sp.; Perkins, 1913 as *Adenoneura falsifalcella*), as well as *Canavalia microcarpa*, *Canavalia tursida*, (and likely other *Canavalia* spp.), and *Strongyloдон ruber* (Fabaceae) (Swezey 1936, 1954, Zimmerman 1978). Swezey ((1908); also quoted in Zimmerman (1978)) provides a detailed description of larval and pupal morphology, and of the biology of this species, paraphrased as follows: eggs are laid singly or in small groups on the seedpod surface; upon hatching larvae bore into the seedpod feeding first on the fleshy walls, then on the ripening seeds; before pupation the larva creates a silken tube to the outer wall of the pod, out of which it eats a hole except for the outermost layer; through which the pupa will protrude to enable adult emergence. Although *Cryptophlebia illepida* can also be bred from *Canavalia* seedpods, the larvae are fairly easily distinguished by the darkly sclerotized thorax of *Cryptophlebia illepida* (Namba 1957, Zimmerman 1978). The ichneumonid wasp, *Trathala flavoorbitalis* Cameron, parasitizes larvae of *C. parapteryx* Swezey (1954). The adults are attracted to lights.

DISTRIBUTION: Endemic to Hawaiian Islands: Oahu – widespread, but only locally common where host plants occur.

REMARKS: Observations of larval biology (Swezey 1908, Perkins 1913) attributed to other *Canavalia*-feeding larvae on Oahu before Meyrick described *C. parapteryx* in 1932, rightly belong to this species (Zimmerman 1978). This is the largest of the Hawaiian species, although adult size varies considerably depending on the quality of the larval diet.

Cydia velocilimitata n. sp.
(Figures 29, 30, 56)

DIAGNOSIS: Most closely allied with *C. falsifalcella*, *C. mauiensis*, and *C. parapteryx*, being the Kauai Island taxon of the *Canavalia*-feeding species group, with wing patterns similar. Differentiated from these other species by its distribution, and aedeagus of male with tip excavated for much shorter distance than in *C. mauiensis*.

DESCRIPTION: (exp. 13-18 mm, n=8) Colored scales on head, body, and legs mostly shades of light ochreous-brown with whitish tips. **Head:** Antennae, head, and labial palpi light brown. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally light brown, tegulae same. Without dorsal tuft of scales. Ventrally lighter. Legs light brown. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figures 29-30) Slightly dilated distally, costa gently arched, apex obtuse, termen somewhat sinuous. Ground color silvery-white. Costal strigulae distinct, plumbaginous, directed towards termen, striae vague below costa in basal area creating a brown-ochreous and white mottled appearance, more distinct and ferruginous distally. Some specimens with a distinct triangular costal patch suffused dark fuscous from base to 2/3 length of costa, posteriorly overlapping the fold. If no costal patch, ferruginous oblique medial fascia extending from end of cell towards basal dorsum, bordered distally by distinct whitish discal patch at end of cell. Ferruginous pretornal patch evident. Ferruginous ocellar area with three distinct ocellar spots, bordered distally by a plumbaginous crescent. Apex with a crescent-shaped ferruginous patch extending into fringe. Continuation of ferruginous stria to central termen extends into fringe, fringe along termen otherwise light ochreous. Venter uniformly light brown-ferruginous. **Hindwings:** Uniformly light brown-ferruginous dorsally, ventrally lighter. Males with glandular ventral patch along path of CuP, opening dorsally and enclosing elongate modified pecten scales. Vein A3 in male displaced towards anal margin. Males without an anal roll or androconial scales (thecae) along vein A3. **Abdomen:** Uniformly light brown. **Male genitalia:** (Figure 56) Tegumen simple, with gently rounded caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa gently concave, ventral edge with deep invagination. Saccus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with dense short setae along ventral and distal margin interspersed with longer setae distally. Aedeagus simple, curved, without cornuti, and with tip somewhat spatulate and excavated dorsally for approximately one-fifth length of aedeagus. **Female genitalia:** (females not known).

TYPE MATERIAL: *Cydia velocilimitata* – ♂ holotype (BPBM): Kauai, Hanalei, Waipa Valley, along Hwy 560, UV light nr. *Canavalia* vines, PT Oboyski, 28.vi.2005, Oboyski specimen PTO-519.2, genitalia slide PTO-s170. 7♂ paratypes (EMEC): Kauai, Hanalei, Waipa Valley, along Hwy 560, UV light nr. *Canavalia* vines, PT Oboyski, 28.vi.2005

ADDITIONAL MATERIAL: This species is known only from a single collection of eight male specimens.

BIOLOGY: Adults were collected at ultraviolet lights in proximity to *Canavalia* vines. Like its close relatives, *C. falsifalcella*, *C. mauiensis*, and *C. parapteryx*, this species probably feeds opportunistically within flower peduncles, stems, and seeds of *Canavalia* spp.

DISTRIBUTION: Endemic to Hawaiian Islands: Kauai. Possibly widespread and locally common where *Canavalia* is abundant, but known only from the type locality at Waipa Valley near Hanalei, Kauai.

REMARKS: Collected along a roadside, where *Canavalia* vines were growing into trees, using an ultraviolet light suspended between a telephone pole and speed limit sign. This type location is the inspiration for the specific epithet.

Cydia plicata (Walsingham 1907)
(Figures 1-4, 15, 17, 31-33, 57, 74)

Adenoneura plicatum Walsingham 1907:678; Plate X, Figure 19 (moth). Swezey & Williams 1932:187, Swezey 1936:198, 1954:204

Cydia plicata. – Zimmerman 1978:609; Figures 374 (wing venation), 382 (moth), 387 (♂ genitalia), 396 (♀ genitalia). Stein 1983a:318, Brenner et al. 2002:101

Cydia crassicornis (sensu Brenner et al. 2002:104, not Walsingham 1907) - misidentification

Cydia falsifalcella (sensu Brenner et al. 2002:104, not Walsingham 1907) - misidentification

Cydia obliqua (sensu Brenner et al. 2002:104, not Walsingham 1907) - misidentification

Cydia storeella (sensu Brenner et al. 2002:104, not Walsingham 1907) - misidentification

DIAGNOSIS: This is a highly polymorphic species in both wing patterns and adult size, and therefore difficult to distinguish from *Cydia makai*, *C. montana*, *C. obliqua*, and *C. storeella*. This species typically is found at higher elevations than *C. makai*, another *Sophora chrysophylla*-feeding species, while *C. montana* is associated with *Acacia koa*. Females can be distinguished from other Hawaiian *Cydia* by the shape of the antrum and lamellae postvaginalis (Figure 74). In males (Figure 57), the tip of the aedeagus is excavated similarly to that of the *Canavalia*-feeding species group and unlike others in the *Sophora*-feeding group.

DESCRIPTION: (exp. 15-20 mm, n=30) Colored scales on head, body, and legs mostly shades of grayish-brown ochreous with whitish tips. **Head:** Antennae, labial palpi, and head light brown-ochreous, head somewhat lighter anteriorly. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally and ventrally light brown ochreous, tegulae same. Without dorsal tuft of scales. Legs light brown ochreous, tibiae somewhat darker. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figures 31-33) Slightly dilated distally, costa gently arched, apex obtuse, termen straight. Ground color ochreous-white, overlaid by highly variable pattern elements. Costal strigulae usually distinct white cinereous but sometimes obscured in specimens with costal suffusion, directed distally towards termen, striae often indistinct in basal half creating mottled appearance or obscured by suffusion of ochreous- to ferruginous-brown triangular costal patch. Oblique medial fascia of ochreous-brown along distal edge of costal triangle extending from end of cell towards basal dorsum sometimes absent. Whitish discal patch usually present and distinct. Pretornal patch of ochreous-brown present or absent, sometimes replaced by whitish scales between dorsal margin and triangular costal patch to ocellar patch. Ocellar patch usually light ochreous-brown grading darker distally with 0, 2, 3 or 4 dark ocellar spots, bordered distally by a silvery-white crescent. Apex often with subtriangular ochreous-brown patch

extending into fringe. Continuation of ochreous-brown stria to central termen extends into fringe. Fringe otherwise light brown-buff. Ventrally uniformly fuscous-brown. **Hindwings:** Dorsally and ventrally light ochreous-brown, darker distally, fringe grayish-white. Males with glandular ventral pouch along path of CuP, opening dorsally and enclosing elongate modified pecten scales. Vein A3 in males displaced towards anal margin. Males without anal roll or androconial scales (thecae) along vein A3. **Abdomen:** Dorsally olivaceous-brown, light brown ventrally. **Male genitalia:** (Figure 57) Tegumen simple with rounded sub-triangular caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa gently concave, ventral edge with deep invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with dense short setae along ventral and distal margin interspersed with longer setae. Aedeagus simple, curved, without cornuti, with somewhat spatulate tip excavated for one-eighth to one-fifth of the length of aedeagus. **Female genitalia:** (Figure 74) Lamellae postvaginalis long and narrow flared into two small lobes caudally. Antrum simple, elongate cylindrical, extending anteriorly into abdominal segment VI with a slight curve. Junction with ductus bursae at anterior end of antrum. Corpus bursae with diverticulum, two long falcate signa. Ductus bursae slightly shorter than width of corpus bursae, with broad tapering junction with corpus bursae to junction with ductus seminalis, narrower from ductus seminalis to antrum.

TYPE MATERIAL: *Adenoneura plicatum* – ♂ holotype (BMNH): Hawaii, Kona, 4000 ft, 13.viii.1892, RCL Perkins; Walsingham specimen 25717; genitalia slide BM 1885. ♀ holotype (BMNH): Hawaii, Kona, over 6000 ft., 30.viii.1892, RCL Perkins; Walsingham specimen 25615. ♂ paratype (BMNH): Hawaii, summit crater of Mt. Hualalai, Kona, 8000 ft, 15.viii.1892, RCL Perkins; Walsingham specimen 28663; F.H. 370. ♂ paratype (BPBM): Hawaii, Mt Kilauea, xii.1896, RCL Perkins; Walsingham specimen 28662; genitalia Busck slide 222; wing slide BM 7570 [there is some confusion about this slide since the specimen still has both pairs of wings]. ♀ allotype (BMNH): Hawaii, Kona, over 6000 ft, 30.viii.1892, RCL Perkins; genitalia slide BM 1886. ♂ homotype (BMNH): Hawaiian Islands, Hawaii, Mauna Loa, 4000 ft, (from seeds of native *Acacia* – [clearly a mistake]), excl. 1900, RCL Perkins; Walsingham specimen 29270; F.H. 679.370.

ADDITIONAL MATERIAL: 2♂, 2♀ (BPBM): Maui, Haleakala, ex: mamane seeds [= *Sophora chrysophylla*], 25.viii.1918, OH Swezey. 1♂, 5♀ (BPBM): Maui, Haleakala, 7000 ft, ex: *Sophora*, 18.viii.1929, OH Swezey; (1♀ abdomen missing). 1♂ (BPBM): Maui, Haleakala, 8000 ft, ex: *Sophora* seeds, 26.ii.1935, OH Swezey. 1 (CAS) HAVO, 5500 ft, 22.v.1986. 3 (UHMA) Hawaii, Mauna Kea. 6♂, 8♀ (EMEC): Hawaiian Islands, Maui, Haleakala N.P., “crater”, 2240 m., reared from *Sophora chrysophylla* seeds, 10.ix.1999, PT Oboyski. 2♀ (EMEC): Hawaiian Islands, Maui, Polipoli, boundary tr., 1830 m., reared from *Sophora chrysophylla* seeds, 31.vii.2002, PT Oboyski. (HAVO) Hundreds of specimens were reared from *Sophora chrysophylla* on Mauna Kea, Hawaii Island between 1995 and 2002 as part of parasitism research (Brenner et al. 2002, Oboyski et al. 2004).

BIOLOGY: Larvae feed within the green (unripe) seeds of *Sophora chrysophylla* (Swezey and Williams 1932, Swezey 1936, 1954, Zimmerman 1978) (Figure 2). Following R.C.L. Perkins’ field notes, Walsingham (1907) mistakenly suggests, “Larva in the seeds of native *Acacias* (Perkins I : 1900).” Eggs are laid singly or in small clusters on the outside of the seedpod (Figure 1), first instar larvae bore into the endosperm. In both the laboratory and the field, larvae may

feed on a single seed, or may tunnel within a seedpod to consume several seeds, and occasionally exit one seedpod and enter another hanging in the same cluster. Occupied seedpods are often detectable by a silk and frass plug covering a hole from the seed to the outside of the pod (Figure 5). Pupation occurs within the seed and the mature pupa protrudes part way out of the silk-covered opening for eclosion of the imago (Figure 4). In the laboratory, some larvae survived for over 18 months in drying seeds and emerged as adults shortly after the application of a moist tissue, suggesting the capacity for facultative diapause. Similarly in the field, I have found live mature larvae in hollowed-out dry mature seeds. Larvae are commonly parasitized by *Pristomerus hawaiiensis*, *Diadegma blackburni*, *Calliephaltes grapholithae* (Ichneumonidae), and *Euderus metallicus* (Eulophidae) (Figure 3) (Brenner et al. 2002, Oboyski et al. 2004). Larvae of this species are also important insect prey of the ‘palila’ (Fringillidae: *Loxioides bailleui*), an endangered Hawaiian honeycreeper bird (Banko et al. 2002). Some aspects of biology are also reviewed by (Swezey 1954, Zimmerman 1978, Oboyski et al. 2004).

DISTRIBUTION: Endemic to Hawaiian Islands: Hawaii – Mauna Kea, Mauna Loa, Hualalai; Maui – Haleakala. Common above 2000 m in *Sophora chrysophylla* forest and shrubland, sympatric with *C. haleakalaensis* and *C. latifemoris* on Maui.

REMARKS: Wing color and patterns are highly variable in this species (Figure 31-33). The high degree of polymorphism resulted in this species being mistakenly identified as several species (*C. crassicornis*, *C. falsifalcella*, *C. obliqua*, and *C. storella*) by Brenner et al. (2002). However, genitalia dissections and DNA sequencing (Oboyski Chapter 3) of specimens obtained by extensive rearing efforts (Brenner et al. 2002, Oboyski et al. 2004) confirm that this is a single variable species. O.H. Swezey (pers. comm. in Zimmerman 1978) notes he has witnessed up to 70% of *Sophora chrysophylla* seeds damaged by *C. plicata* larvae. However, *C. haleakalaensis*, *C. latifemoris* and *C. makai* also feed on *S. chrysophylla* seeds and may be partly responsible for Swezey’s observation. The adult moths are attracted to ultraviolet lights.

Cydia obliqua (Walsingham 1907)
(Figure 75)

Enarmonia (?) *obliqua* Walsingham 1907:686; Plate XI. Figure 4 (moth)

Adenoneura obliqua. – Meyrick 1932:222

Cydia obliqua. – Zimmerman 1978:608; Figures 381 (moth), 394 (♀ genitalia)

Cydia oblique. – Stein 1983a:318 (misspelling of *C. obliqua* Walsingham)

DIAGNOSIS: *Cydia obliqua*, known from only three female specimens, closely resembles a form of the polymorphic *C. plicata* found on Hawaii and Maui. However, the short, stout, female antrum and wide lamellae postvaginalis of *C. obliqua* (Figure 75) is distinctly different from other Hawaiian *Cydia*.

DESCRIPTION: (exp. 14-17 mm, n=3) **Head:** Antennal scales fuscous with lighter tips, labial palpi and head buff white. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsal scale light brown-cinereous with lighter tips, mid-dorsum somewhat darker, tegulae light brown-cinereous becoming lighter with longer scales posteriorly. Without dorsal tuft of scales. Ventral body and

legs light brownish-buff. No discernable sex scales (e.g. hair pencils). **Forewings:** Slightly dilated distally, costa gently arched, apex obtuse, termen nearly straight. Ground color whitish-cinereous. Costal strigulae somewhat obscured basally, distally distinct directed distally towards termen, striae indistinct in basal half. Mid-basal area buff white. No triangular costal patch. Oblique medial fascia light olivaceous-brown extending from end of cell towards basal dorsum ending in a darker dorsal blotch. Whitish discal patch distinct. Pretornal patch light olivaceous-brown basally, darker distally. Ocellar patch light brown-cinereous with two or three indistinct ocellar spots, bordered distally by a silvery-brown crescent. Apex with a light olivaceous-brown patch extending into fringe. Continuation of olivaceous-brown stria to central termen extends into fringe. Fringe otherwise buff white. Ventrally uniform dark brown-cinereous, ventral fringe matching dorsum. **Hindwings:** Dorsally brown-cinereous, darker distally, buff gray costal margin, fringe buff gray. Ventrally similar to dorsum but somewhat lighter. It is not known whether males have a glandular ventral pouch. **Abdomen:** Light brown-cinereous dorsally, lighter ventrally. **Male genitalia:** (males not known). **Female genitalia:** (Figure 75) Lamellae postvaginalis somewhat wider than long, dilated posteriorly. Antrum funnel-shaped with two lightly sclerotized longitudinal bands, direct anterior junction with ductus bursae. Corpus bursae with diverticulum, two long falcate signa. Ductus bursae somewhat shorter than width of corpus bursae, wider between corpus bursae and ductus seminalis than between ductus seminalis and antrum.

TYPE MATERIAL: *Enarmonia (?) obliqua* – ♀ holotype (BMNH): Hawaii, Hualalai (Kona), 5000 ft., 15.VIII.1892, RCL Perkins; Walsingham specimen 25828; genitalia slide BM 1882. ♀ paratype (BMNH): Hawaii, Hualalai (Kona), 5000 ft., 15.VIII.1892, RCL Perkins; Walsingham specimen 25832.

ADDITIONAL MATERIAL: 1♀ (BMNH): Hawaii, Kona, 4000 ft., 10.VIII.1892, RCL Perkins; Walsingham specimen 25271; (with determination label: *Enarmonia* sp. Drnt.).

BIOLOGY: There is no information available regarding larval biology, host plants, habitat, predators, or parasitoids.

DISTRIBUTION: Endemic to Hawaiian Islands: Hawaii – Known only from three female specimens collected 1200-1500 m on Hualalai volcano.

REMARKS: If not an extinct species (or nearly so), it is possible that the three specimens of *C. obliqua* represent an aberrant form of the highly polymorphic *C. plicata* found commonly at higher elevations (> 2000 m) on Hawaii and Maui. But without further material showing greater similarity to the type specimens I shall resist synonymizing these two species. Specimens misidentified as *C. obliqua* by Brenner et al. (2002) are correctly referred to *C. plicata*.

Cydia storeella (Walsingham 1907)
(Figure 77)

Enarmonia (?) *storeella* Walsingham 1907:686; Plate XI. Figure 3 (moth)

Adenoneura storeella. – Meyrick 1932:222

Cydia storeella. – Zimmerman 1978:609; Figures 383 (moth), 398 (♀ genitalia)

DIAGNOSIS: Color and pattern of wings similar to morphs of *Cydia plicata* (e.g. Figure 33). Distinguished from all other female Hawaiian *Cydia* by the elongate antrum and shape of the lamellae postvaginalis. Males not known.

DESCRIPTION: (exp. 14 mm, n=1) Colored scales on head, body, and legs mostly shades of brown-cinereous with whitish tips. **Head:** Antennae, head, and labial palpi light brown-cinereous, head somewhat lighter anteriorly. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally light brown-cinereous, tegulae same. Ventrally whitish-buff (although largely denuded of scales). Without dorsal tuft of scales. Legs light brown-cinereous. No discernable sex scales (e.g. hair pencils). **Forewings:** Slightly dilated distally, costa gently arched, apex obtuse, termen straight. Ground color whitish-buff. Costal strigulae not apparent. Distinct triangular costal patch suffused brown-olivaceous from base to 2/3 length of costa, posteriorly overlapping fold, bordered distally by a lighter band of scales and distinct whitish discal patch at end of cell. Pretornal patch only vaguely evident as light brown-cinereous. Ocellar patch indistinct and with no discernable ocellar spots or distal border, and no discernable apical patch. Distal area instead speckled light brown olivaceous, with thin brown-olivaceous line along termen. Fringe uniformly light plumbaginous. Ventrally light brown-cinereous. **Hindwings:** Uniformly brown-cinereous dorsally, ventrally lighter. It is unknown whether males of this species possess a ventral pouch. **Abdomen:** (removed). **Male genitalia:** (males not known). **Female genitalia:** (Figure 77) Lamellae postvaginalis slightly longer than wide, constricted centrally and widely flared posteriorly. Antrum very long and slender relative to other Hawaiian *Cydia*, projecting with a lightly sclerotized ventral lobe just beyond junction with ductus bursae. Corpus bursae with diverticulum. Two long, falcate signa on opposite sides of corpus bursae (not proximate as suggested by Zimmerman 1978). Ductus bursae shorter than width of corpus bursae, sharply tapering from corpus bursae, then uniformly slender to junction with antrum.

TYPE MATERIAL: *Enarmonia* (?) *storeella* – ♀ holotype (BMNH): Hawaiian Islands, Maui, Haleakala, 5000 ft, v.1896, RCL Perkins; Walsingham specimen 28185; genitalia slide BM 1881.

ADDITIONAL MATERIAL: This species is known only from the female holotype.

BIOLOGY: There is no information available regarding larval biology, host plants, habitat, predators, or parasitoids.

DISTRIBUTION: Endemic to Hawaiian Islands: Maui – Haleakala, 5000 ft. Known from a single female specimen.

REMARKS: If not extinct, this species is very rare. There remains ample native forest around 5000 ft on Haleakala, Maui, so it is doubtful that this species was lost due to habitat destruction, although many species of forest understory plants have become rare or extirpated due to ungulate and rat feeding. Zimmerman (1978:607) suggested the approximation of the signa in the corpus bursae is a diagnostic characteristic for this species. However, closer examination of the holotype slide (BM 1881) reveals that the signa are on opposite sides of the corpus bursae giving the appearance of proximity. The antrum of the female genitalia appears unique and not likely to be an aberrant form of another more common polymorphic species (e.g. *C. plicata*). However, *C. crassicornis* (Hawaii) and *C. gypsograptia* (Oahu) are known from only male holotypes. Specimens misidentified as *C. storeella* by Brenner et al. (2002) are correctly referred to *C. plicata*.

Cydia haleakalaensis n. sp.
(Figures 34, 58, 77)

DIAGNOSIS: This species is uniformly dark (mottled dark brown and charcoal), which is unique among Hawaiian *Cydia*.

DESCRIPTION: (exp. 12-17 mm, n=6) Colored scales on head, body, and legs fuscous with whitish tips. **Head:** Head scales fuscous tipped with silvery-white giving a frosted appearance. Labial palpi pale cinereous, lighter towards tips. Antennae banded fuscous and pale cinereous. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsal color same as head, tegulae same. Ventrally similar. Without dorsal tuft of scales. Legs slightly darker. No discernable sex scale (e.g. hair pencils). **Forewings:** (Figure 34) Slightly dilated distally, costa gently arched, apex obtuse, termen nearly straight. Ground color fuscous. Costal strigulae vague, fuscous tipped with light cinereous, interspersed with dark brown-ferruginous streaks, directed towards termen, striae indistinct giving a mottled appearance throughout. No triangular costal patch. Oblique medial fascia of dark brown-ferruginous extending from end of cell towards basal dorsum. Discal patch absent. Dark brown-ferruginous pretornal patch vaguely present. Ocellar patch indistinct and without ocellar spots. Termen, from apex to tornus, with a band of brown-ferruginous mottled with fuscous. Fringe at apex light brown-ferruginous, dark brown fuscous below apex. Ventrally uniform dark-brown fuscous. **Hindwings:** Dorsal and ventral dark brown fuscous. Males with glandular ventral pouch along path of CuP, opening dorsally and enclosing elongate modified pecten scales. Vein A3 in males displaced towards anal margin. Males without anal roll or androconial scales (thecae) along vein A3. **Abdomen:** Dark brown-fuscous. **Male genitalia:** (Figure 58) Tegumen simple with central point between short lobes that create a somewhat parallel-sided crista along caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa gently concave, ventral edge with deep invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with dense short setae along ventral and distal margin interspersed with longer setae distally. Aedeagus simple, curved, without cornuti. Tip of aedeagus with two dorsal ridges and often a short, sharp ventral projection. **Female genitalia:** (Figure 77) Lamellae postvaginalis longer than wide, constricted through middle, and flared posteriorly. Antrum forms a stout cylinder as wide as ostial opening directly into ductus bursae. Corpus bursae with diverticulum, two long falcate signa. Ductus bursae longer than width of corpus, tapering gradually from corpus bursae to antrum.

TYPE MATERIAL: *Cydia haleakalaensis* - ♂ holotype (BPBM) Maui, Haleakala N.P. nr. Hosmer grove, 2105 m., aerial netting, 1.vii.2005, PT Oboyski. Specimen PTO-521.1, genitalia slide pto-s146. ♀ allotype (BPBM) Maui, Haleakala N.P., nr. Oilipuu, 2040 m., reared: *Sophora chrysophylla* pods, 8.vii.2005, PT Oboyski. Specimens PTO-538.22, genitalia slide pto-s337. 3♂ paratypes (EMEC) Maui, Haleakala N.P. nr. Hosmer grove, 2105 m., aerial netting, 1.vii.2005, PT Oboyski. Specimens PTO-521.2,3,4.

ADDITIONAL MATERIAL: 1♂ (EMEC) Maui, Haleakala N.P., nr. Hosmer grove, 2105 m., reared: *Sophora chrysophylla* pods, 29.vii.2002, PT Oboyski. Specimen greasy. 1♀ (EMEC) Maui, Haleakala N.P., Kaupo Gap, 1591 m., reared: *Sophora chrysophylla* pods, 5.viii.2002, PT Oboyski. Specimen greasy. 5♂, 1♀ (EMEC) Maui, Haleakala N.P., Halemauu Trail, 2240 m., reared: *Sophora chrysophylla* pods, 6.viii.2002, PT Oboyski. Specimens greasy.

BIOLOGY: Larvae feed within the ripening seeds of *Sophora chrysophylla*. Four adult male specimens of this species were captured flying during the day. It is unclear whether this is a diurnal species, or if these individuals were flushed from the vegetation. However, each was observed from some distance, suggesting they were not flushed, and none has been collected at lights.

DISTRIBUTION: Endemic to Hawaiian Islands: Maui – Haleakala, 1900-2240 m., along the outer slopes and within the summit “crater,” sympatric with *C. latifemoris* and *C. plicata*.

REMARKS: Molecular evidence places *C. haleakalaensis* sister to *C. latifemoris* (Oboyski Chapter 3), with which it is sympatric.

Cydia latifemoris (Walsingham 1907)
(Figures 35, 59, 78)

Adenoneura latifemoris Walsingham 1907:679; Plate X. Figure 20 (moth). Swezey 1936:198, 1954: 204

Cydia latifemoris. – Zimmerman 1978:592; Figures 373 (wing venation), 380 (moth), 385 (♂ genitalia), 392 (♀ genitalia)

Cydia latefemoris. – Miller 1990:126 (misspelling of *latifemoris*)

DIAGNOSIS: The rich, dark brown and black coloring of fresh specimens of this species and *C. haleakalaensis* are distinct among Hawaiian *Cydia*. This species can be distinguished from its sympatric sister-species, *C. haleakalaensis* (uniformly dark), by its distinct pretornal blotch (Figure 35), the excavated tip of the male aedeagus (Figure 59), and the more distinct ventral lobe of the female antrum (Figure 78).

DESCRIPTION: (exp. 11-16, n=10) Colored scales on head, body, and legs mostly shades of dusky-brown with whitish tips. **Head:** Antennae, head and labial palpi dark brown-cinereous. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally dark brown-cinereous, tegulae same. Without dorsal tuft of scales. Ventrally lighter, somewhat ochreous around coxae. Legs light

brown-cinereous. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figure 35) Slightly dilated distally, costa gently arched, apex obtuse, termen nearly straight. Ground color light brown-cinereous. Costal strigulae distinct, directed distally towards termen, striae indistinct in basal half creating mottled appearance. No triangular costal patch. Oblique medial fascia extending from end of cell towards basal dorsum. Whitish discal patch present but indistinct. Pretornal patch light brown-ochreous basally with distal margin dark brown-cinereous. Ocellar patch light brown-ochreous with three distinct ocellar spots, bordered distally by a silvery crescent. Apex with a light brown-ochreous patch with dark brown-cinereous distal edge. Continuation of light brown-ochreous stria streaked dark brown-cinereous to central termen along termen edge to tornus, bordered distally by dark brown-cinereous. Fringe silvery-bronze. Ventrally uniformly light brown-cinereous, lighter along anal margin. **Hindwings:** Dorsal and ventral dark brown-cinereous distally grading to lighter brown basally. Males with glandular ventral pouch along the path of CuP opening dorsally and enclosing elongate modified pecten scales. Vein A3 in males displaced towards anal margin. Males without anal roll or androconial scales (thecae) along vein A3. **Abdomen:** Uniformly dark brown-cinereous. **Male genitalia:** (Figure 59) Tegumen simple with single-lobed crista along caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa gently concave, ventral margin with a deep invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with dense short setae along ventral and distal margin. Aedeagus simple, curved, without cornuti, dilated distally with somewhat spatulate tip excavated dorsally for one-eighth of the length of aedeagus. **Female genitalia:** (Figure 78) Lamellae postvaginalis somewhat longer than wide, tightly constricted centrally and wider posteriorly than at ostium. Antrum elongate cylindrical with ventral lobe extending beyond junction with ductus bursae. Corpus bursae with diverticulum, two long falcate signa. Ductus bursae slightly shorter than width of corpus bursae, only slightly wider at junction with corpus bursae than junction with antrum.

TYPE MATERIAL: *Adenoneura latifemoris* – ♀ holotype (BMNH): Hawaiian Islands, Maui, Haleakala crater, X.1896, RCL Perkins; Walsingham specimen 28127; genitalia slide BM 2054. ♂ allotype (BMNH): Hawaii: Hualalai (Kona), summit of crater, 8000 ft., 15.VII.1892, RCL Perkins; Walsingham collection 28665; genitalia slide BM 2053. ♀ paratype (BPBM): Hawaii: Hualalai (Kona), summit of crater, 8000 ft., 15.VII.1892, RCL Perkins; Walsingham collection 28664; (abdomen missing); “Both sexes occurred at the summit of Hualalai but the ♀ is in poor condition” and therefore used as a paratype rather than the holotype (Walsingham 1907:679).

ADDITIONAL MATERIAL: 3♂ (BPBM): Maui, Haleakala, ex: mamane seeds [= *Sophora chrysophylla*], 25.viii.1918, OH Swezey; missing parts – (1♂ abdomen, antennae, hindwing sexpouchs; 1♂ head; 1♂ left hindwing sexpouch). 1♀ (BPBM): Maui, Haleakala, ex: mamane seeds [= *Sophora chrysophylla*], 25.viii.1918, OH Swezey; (abdomen missing). 2♂ (BPBM): Maui, Haleakala, 8000 ft, ex: *Sophora* seeds, 26.ii.1935, OH Swezey. 5♀ (EMEC): Hawaiian Islands, Maui, Haleakala N.P., “crater”, 2240 m., reared from *Sophora chrysophylla* seeds, 10.ix.1999, PT Oboyski. 1♀ (EMEC): Hawaiian Islands, Maui, Polipoli, boundary tr., 1830 m., reared from *Sophora chrysophylla* seeds, 31.vii.2002, PT Oboyski.

BIOLOGY: Larvae feed within the ripening seeds of *Sophora chrysophylla* (Swezey 1936, 1954, Zimmerman 1978) at mid to high elevations (> 2000 m). O.H. Swezey (pers. comm. in Zimmerman, 1978) noted that the larvae of *C. latifemoris* can destroy nearly half of the seed crop

of *Sophora*. On Maui (Haleakala), *C. latifemoris* is sympatric with *C. plicata* and *C. haleakalaensis*, which also feed in the seeds of *Sophora*. It is unclear, therefore, if Swezey was referring specifically to *C. latifemoris*. I also observed terminal twigs of *Sophora chrysophylla* with past feeding damage (*i.e.* split in terminal twigs) similar to those on Kauai from which *C. makai* were reared. However, no larvae have been found in twigs on Maui. Adults were collected after sunset at ultraviolet lights (compare with *C. haleakalaensis*). Larvae are parasitized by the ichneumonid wasps *Pristomerus hawaiiensis* Perkins, *Diadegma blackburni* Cameron, and *Calliephialtes grapholithae* Cresson, as well as the eulophid wasp *Euderus metallicus* Ashmead.

DISTRIBUTION: Endemic to Hawaiian Islands: Maui – *C. latifemoris* is particularly abundant in the *Sophora* shrublands of Haleakala volcano, both on the outer slopes and within the “crater,” sympatric with *C. haleakalaensis* and *C. plicata*. Despite targeted collection efforts, no other specimens of this species have been collected on Hawaii Island since the two specimens collected by Perkins in 1892. This species is probably not a persistent resident of Hawaii Island. However, given the proximity (~100 km) of Haleakala and Hualalai it would not be surprising for *C. latifemoris* to periodically establish small populations in the subalpine *Sophora* forests of Hualalai.

REMARKS: Specimens misidentified as *C. latifemoris* by Brenner et al. (2002) are correctly referred to *C. plicata*.

***Cydia makai* n. sp.**
(Figures 36, 37, 60, 79)

DIAGNOSIS: *Cydia makai* is polymorphic. Forms of *C. makai* are nearly identical to forms of the highly polymorphic *C. plicata*. A slight difference in the shape of the tip of the male aedeagus can be used to separate these species. *Cydia makai* is found at elevations lower than other *Sophora*-feeding species (<2000 m), although *C. makai* distributions overlap with *C. plicata* on Hawaii Island. Regarding molecular data, *C. makai* and *C. plicata* are consistently divergent for the mitochondrial genes cytochrome oxidase I and II (2.9-4.3% and 2.3-3.8% uncorrected P, respectively) when comparing specimens pooled across islands or within individual islands separately (Oboyski Chapter 3).

DESCRIPTION: (exp. 11-17 mm, n=8) Colored scales on head, body, and legs mostly shades of light brown-ferruginous with whitish tips. **Head:** Antennae, head, and labial palpi light brown-ferruginous. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Similar color to head, tegulae tending more ferruginous posteriorly. Without dorsal tuft of scales. Legs light brown-ferruginous. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figures 36-37) Slightly dilated distally, costa gently arched, apex obtuse, termen straight. Ground color light brown-buff. Most commonly (Figure 36), costal strigulae distinct, light brown-ferruginous, basally interspersed with dark ferruginous becoming lighter distally, directed towards termen, striae indistinct except near apex. Basal area light brown-buff, mottled with light brown-ferruginous. No triangular costal patch, oblique medial fascia, discal patch or pretornal patch. Ocellar patch vaguely defined by light brown ferruginous with three somewhat distinct ocellar spots, bordered distally by a plumbaginous crescent. Apical patch indistinct. Termen below apex to tornus with a

ferruginous band. Fringe ferruginous, becoming fuliginous towards tornus. An alternative form on Kauai (Figure 37) with ferruginous triangular costal patch bordered distally by light brown-ferruginous. A dark fuscous streak runs through the cell. Termen, from apex through tornus, bordered by dark fuscous band. Fringe ferruginous mixed with fuscous. Ventrally uniform light brown-ferruginous. **Hindwings:** Dorsal and ventral light bronze-ferruginous, somewhat lighter basally and ventrally. Males with glandular ventral pouch along path of CuP, opening dorsally and enclosing elongate modified pecten scales. Vein A3 in males displaced towards anal margin. Males without anal roll or androconial scales (thecae) along vein A3. **Abdomen:** light brown-ferruginous dorsally, lighter ventrally. **Male genitalia:** (Figure 60) Tegumen simple with rounded caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa gently concave, ventral edge with deep invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with dense short setae along ventral and distal margin densely intersperse with longer setae. Aedeagus simple, curved, without cornuti, and with tip excavated dorsally approximately one-eighth the length of aedeagus. **Female genitalia:** (Figure 79) Lamellae postvaginalis long and narrow flared into two small lobes caudally. Antrum simple, elongate cylindrical, extending anteriorly into abdominal segment VI with a slight dorsal curve. Junction with ductus bursae at anterior end of antrum. Corpus bursae with diverticulum, tow long falcate signa. Ductus bursae slightly shorter than width of corpus bursae, with broad tapering junction with corpus bursae to junction of ductus seminalis, narrower from ductus seminalis to antrum.

TYPE MATERIAL: *Cydia makai* – ♂ holotype (BPBM): Hawaii Islands, Molokai, TNC Kamakou Preserve, nr. Onini gulch, northern branch, 902 m., 25.vii.2003, PT Oboyski. Specimen PTO-197.7. ♀ holotype (BPBM): Hawaii Islands, Molokai, TNC Kamakou Preserve, nr. Onini gulch, northern branch, 902 m., 25.vii.2003, PT Oboyski. Specimen PTO-197.56. 1♂ paratype (EMEC) Hawaii Islands, Molokai, TNC Kamakou Preserve, nr. Onini gulch, southern branch, 881 m., 25.vii.2003, PT Oboyski. Specimen PTO-197.11, genitalia slide pto-s311. 1♂ paratype (EMEC) Hawaii Islands, Molokai, TNC Kamakou Preserve, nr. Onini gulch, southern branch, 881 m., 25.vii.2003, PT Oboyski. Specimen PTO-198.21, genitalia slide pto-s148. 2♀ paratypes (EMEC) Hawaii Islands, Molokai, TNC Kamakou Preserve, nr. Onini gulch, northern branch, 902 m., 25.vii.2003, PT Oboyski. Specimen PTO-197.25, genitalia slide pto-s147; specimen PTO-197.31, genitalia slide pto-s152.

ADDITIONAL MATERIAL: 1♀ (EMEC): Hawaiian Islands, Kauai, Kokee State Park, Awaawapuhi trail, 1101 m, at UV light, 26.ii.2005, PT Oboyski. 1♂ (EMEC): Hawaiian Islands, Kauai, Kokee State Park, Nualolo trail, 1049 m., reared from *Sophora chrysophylla* live terminal twig, 27.ii.2005, PT Oboyski. 2♂, 1♀ (EMEC): Hawaiian Islands, Hawaii, Kohala, Kawaihae Uka, *Acacia koaia* sactuary, 965 m., reared from *Sophora chrysophylla* seeds, 5.vii.2003, PT Oboyski. 2♂, 2♀ (EMEC): Hawaiian Islands, Hawaii, Kilauea, HAVO N.P., Ainahou, 930 m., reared from *Sophora chrysophylla* seeds, 6.vii.2003, PT Oboyski.

BIOLOGY: Larvae feed within the ripening seeds of *Sophora chrysophylla*, and are parasitized by *Calliephialtes grapholithae*, *Diadegma blackburni*, *Pristomerus hawaiiensis*, and *Euderus metallicus*. A non-native species of *Araecerus* (Anthribidae) beetle also feeds within seeds of *Sophora* at low elevations. It is unclear if *Cydia makai* compete directly with beetles, which are

more often found in mature, hardened seeds. However, very few intact seeds are added to the local seed bank in areas with beetle infestation (PT Oboyski, personal observation).

DISTRIBUTION: Endemic to Hawaiian Islands: low elevations (< 2000 m) on Kauai, Molokai, and Hawaii, and probably occurred on Oahu, Lanai, and Maui. Bishop Museum herbarium specimens of *Sophora chrysophylla* seedpods from Oahu and Lanai have exit holes similar to those made by *Cydia* species.

REMARKS: The specific epithet *makai* in the Hawaiian language means towards the sea, as opposed to *mauka*, meaning towards the mountains. This species is found at elevations lower than other *Sophora*-feeding species, and is difficult to identify where their ranges overlap. *Cydia makai* will likely go extinct within the next 100 years as the number of low elevation *Sophora* trees continues to decline, particularly on the older islands, and as the alien seed beetle, *Araecerus* sp., expands its range.

Cydia koaiae n. sp.
(Figures 10, 38, 39, 63)

Cydia new species 1 Zimmerman 1978:610; Figures 377 (feeding damage), 386 (♂ genitalia)

DIAGNOSIS: *Cydia koaiae*, endemic to Hawaii Island, is a gray and black species likely to be confused with its sister-species, *C. conspicua*, known only from the older islands. For most male specimens of *C. koaiae* the cucullus has an abrupt angle not present in *C. conspicua*.

DESCRIPTION: (exp. 11-21 mm, n=12) Scale colors on head, body, and legs mostly shades of grayish-brown with whitish tips. **Head:** Antennae, head, and labial palpi uniformly light brown-cinereous, head and palpi somewhat lighter on top. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally and ventrally light brown-cinereous, lighter ventrally, tegulae the same. Without dorsal tuft of scales. Legs whitish-fuscous, fore- and middle-legs tibia and tarsomeres banded with fuscous bases and lighter tips. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figures 10, 38, 39) Slightly dilated distally, costa gently arched, apex obtuse, termen somewhat sinuate. Ground color light cinereous. Costal strigulae distinct, directed distally towards termen, but somewhat obscured when triangular costal patch is present. Distinct triangular costal patch present or absent. If present, fuliginous-brown suffused from base to two-thirds length of costa, posteriorly overlapping the fold. If absent, basal area streaked with striae, and oblique medial fascia of fuliginous-brown extending from end of cell towards basal dorsum. Distal end of cell with a conspicuous white discal patch. Costal patch with a fuliginous-brown projection from below discal patch connecting to a vague pretornal blotch. Ocellar patch light cinereous with three distinct ocellar spots, bordered distally by a silvery-white crescent. Apex with a subtriangular fuliginous-brown patch extending into fringe. Continuation of fuliginous-brown stria to central termen extends into fringe. Fringe otherwise silvery-white. Ventrally light fuliginous-brown. **Hindwings:** Dorsally light brown-cinereous, somewhat lighter basally and ventrally. Males with a ventral glandular pouch below cubital vein opening dorsally and enclosing elongate modified pecten scales. Vein A3 in males displaced towards anal margin. Males without an anal roll or androconial scales (thecae) along vein A3. **Abdomen:** Mottle

brown-cinereous and cinereous, somewhat lighter ventrally. **Male genitalia:** (Figure 63) Tegumen simple, lacking gnathos, socii, and uncus, with crista of caudal ridge broadly rounded with central point. Valvae with excavation in basal third, costa gently concave, ending at an abrupt angle, ventral edge with deep invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate, rounded below abrupt angle at end of costal edge, with dense short and long setae along ventral and distal margin. Aedeagus simple, curved, without cornuti, spatulate tip excavated dorsally for one-fifth length of aedeagus. **Female genitalia:** (females not known).

TYPE MATERIAL: *Cydia koaiae* – ♂ holotype (BPBM): Hawaiian Islands, Hawaii, Kohala, Kawaihae Uka, Acacia koaia sanctuary, 970 m., at UV light, 18.vii.2004, PT Oboyski; Oboyski specimen PTO-323.13, genitalia slide pto-s333. 11♂ paratypes (EMEC): Hawaiian Islands, Hawaii, Kohala, Kawaihae Uka, Acacia koaia sanctuary, 970 m., at UV light, 18.vii.2004, PT Oboyski.

ADDITIONAL MATERIAL: 2♂ (HDOA): Hawaii, Kawaihae Uka, ex: *Acacia koaia* dead twig, 3.iii.52, C.J. Davis collector. 3♂ (HDOA): Hawaii, Kawaihae Uka, ex: *koaia*, 17.ii.65, C.J. Davis collector, H-65-1. 1♂ (HDOA): Hawaii, Kawaihae Uka, ex: *Acacia koaia*, 30.i.68, C.J. Davis collector.

BIOLOGY: The larvae of *C. koaiae* feed in the senescing terminal branches of *Acacia koaia*, along with *C. walsinghamii*. It is unclear whether larvae cause the senescence or infest already senescing branches. Larvae are most likely attacked by the suite of *Cydia* parasitoids in Hawaii, although I know of none that has been reared. Host trees at the *Acacia koaia* sanctuary on Hawaii Island are interspersed with *Sophora chrysophylla* from which I have reared *Cydia makai* and its parasitoids, *Pristomerus hawaiiensis*, *Diadegma blackburni*, *Calliephialtes grapholithae* (Ichneumonidae), and *Euderus metallicus* (Eulophidae). The adults of *C. koaiae* are attracted to lights.

DISTRIBUTION: Endemic to Hawaiian Islands: Hawaii – Kohala, 950-1000 m. Appears to be limited to a small, protected population of *Acacia koaia* trees in the Kohala district of Hawaii Island. Although a small population of *Acacia koaia* exists on Molokai, a single *Cydia* specimen reared from these trees is clearly *C. walsinghamii*. *Acacia koaia* also grows on Maui, but I have not seen these trees, nor do I know of any moths that have been collected in association with them.

REMARKS: Zimmerman (1978) qualified this as a new species that may have been confused with *C. walsinghamii* by past researchers, but declined to name it. Unlike *C. walsinghamii*, however, *C. koaiae* has the sex pouch in the male hindwing. This species was discovered by the late entomologist C.J. Davis in 1965, who first investigated the senescing branches of *Acacia koaia* at its sanctuary enclosure in Kohala, Hawaii (see photo in Zimmerman 1978: figure 377).

Cydia conspicua (Walsingham 1907)
(Figures 40, 41, 64, 82)

Enarmonia (?) *conspicua* Walsingham 1907:684; Plate X. Figure 28 (moth)

Adenoneura conspicua. – Meyrick 1928:98; 1932: 222. Swezey 1954:4

Cydia conspicua. – Zimmerman 1978:585; Figures 372 (wing venation), 376 (pupa), 378 (moth), 385 (♂ genitalia), 390-391 (♀ genitalia)

DIAGNOSIS: Most closely allied with *C. koaiae* from Hawaii Island, sharing similar black, gray, and white color patterns on the forewings. Males of *C. conspicua* with cucullus elongate rounded while most males of *C. koaia* have an angle near the apex of cucullus. Alternate forms of *C. conspicua* on Oahu (Figure 41) more closely resemble the color patterns of *C. walsinghamii*, but males of *C. walsinghamii* lack the hindwing pouch.

DESCRIPTION: (exp. 12-20 mm, n=27) Scale colors on head, body, and legs mostly shades of grayish-brown with whitish tips. **Head:** Antennae and labial palpi uniformly brownish-cinereous, palpi somewhat lighter on top. Head uniformly whitish-brown. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally and ventrally brownish-cinereous, lighter ventrally, tegulae somewhat ferruginous. Without dorsal tuft of scales. Legs whitish-fuscous, tibia darker than femur, tarsomeres banded with fuscous bases and lighter tips. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figures 40-41) Slightly dilated distally, costa gently arched, apex obtuse, termen somewhat sinuate. Ground color ochreous-white, overlaid by variable pattern elements. Costal strigulae distinct, directed distally towards termen. Most often with a distinct triangular costal patch suffused from base to two-thirds length of costa, posteriorly overlapping the fold, ranging in color from fuscous to ferruginous. Costal patch sometimes not fully suffused anterior-basal leaving an indistinct basal suffusion or an oblique medial fascia extending from end of the cell towards basal dorsum. Distal end of cell with a conspicuous white discal patch. Occasionally costal patch with a fuscous projection from below discal patch towards tornus (as in ♀ holotype) or a triangular fuscous pretornal blotch. Pale ocellar area with three or four ocellar spots not always visible, bordered distally by a lighter brownish-white crescent. Apex with a subtriangular fuscous-ferruginous patch extending into fringe. Continuation of fuscous-ferruginous stria to central termen extends into fringe. Ventrally brown-fuscous with pale basal anal area. **Hindwings:** Uniformly brown-fuscous, somewhat paler ventrally. Males with glandular ventral pouch along path of CuP opening dorsally and enclosing elongate modified pecten scales. Vein A3 in males displaced towards the anal margin. Males without an anal roll or androconial scales (thecae) along vein A3. **Abdomen:** Uniformly fuscous. **Male genitalia:** (Figure 64) Tegumen simple, with bilobed or rounded crista along the caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa slightly concave, ventral margin with shallow invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with dense short setae along ventral and distal margin, interspersed with fewer long setae. Aedeagus simple, strongly curved, without cornuti, spatulate tip excavated dorsally for one-fifth length of aedeagus. **Female genitalia:** (Figure 82) Lamellae postvaginalis slightly longer than wide, hourglass-shaped. Antrum without complex sclerotization pattern, elongate with a lobe extending just beyond junction with ductus bursae. Corpus bursae with diverticulum. Two long, falcate signa on opposite side of corpus bursae (no proximate as suggested by

Zimmerman 1978). Ductus bursae short, joined broadly to corpus bursae, quickly tapering to antrum.

TYPE MATERIAL: *Enarmonia (?) conspicua* Walsingham – ♀ holotype (BMNH): Hawaiian Islands, Maui, Haleakala, 5000 ft., x.1896, RCL Perkins; Walsingham specimen 28134; genitalia slide BM 2052. ♂ paratype (BMNH): Oahu, Honolulu, P . 08, RCL Perkins. 2♂, 2♀ paratypes (BMNH): Oahu, Kahauiki, bred [*Acacia koa*], OH Swezey, 28.ix.1924; [includes 2♂ pupal cases, 1♀ head in a vial].

ADDITIONAL MATERIAL: 1♂ (BPBM): Oahu, Sugarloaf hill, ex: koa seeds [= *Acacia koa*], 15.xi.1922, OH Swezey. 3♀ (HDOA): Oahu, Kahauiki, ex: koa bark, 28.ix.1924, OH Swezey; (determined by Meyrick). 4 (HDOA): Oahu, Kahauiki, ex: koa, 14.ix.1930, OH Swezey. 2♂ (HDOA): Oahu, Tantalus, 25.x.1930, FC Hadden; (determined by Meyrick). 1♀ (EMEC): Hawaiian Islands, Kauai, Kokee State Park, Canyon trail, trail head, 1052 m., at UV light, 28.ii.2005, PT Oboyski. 2♂ (EMEC): Hawaiian Islands, Kauai, Kokee State Park, Canyon trail, trail head, 1052 m., at UV light, 5.iii.2005, PT Oboyski. 3♂ (EMEC): Hawaiian Islands, Kauai, Kokee State Park, Makaha road, 1040 m., at UV light, 17.vi.2005, PT Oboyski. 1♂, 1♀ (EMEC): Hawaiian Islands, Kauai, Kokee State Park, Kumuwela ridge, 1113 m., at UV light, 20.vi.2005, PT Oboyski. 1M (EMEC): 1♂, 1♀ (EMEC): Hawaiian Islands, Kauai, Kokee State Park, Awaawapuhi trail, 1027 m., at UV light, 21.vi.2005, PT Oboyski. 3♂ (EMEC): Hawaiian Islands, Kauai, Kokee State Park, Awaawapuhi trail, 1171 m., at UV light, 23.vi.2005, PT Oboyski. 1♂ (EMEC): Hawaiian Islands, Kauai, Kokee State Park, 4WD road off Canyon road, 887 m., at UV light, 26.vi.2005, PT Oboyski. 1♂, 3♀ (EMEC): Hawaiian Islands, Oahu, S. Waianae Mtns., TNC Honouliuli preserve, Kaluaa section, 607 m., at UV light, 22.vii.2005, PT Oboyski. 1♂, 1♀ (UHMA): HI: Oahu, Koolau Mtns, Wiliwilinui Trail, UV light trap, 6 June 2006, J. Eiben & W. Haines.

BIOLOGY: Larvae feed in the seeds and decaying bark of *Acacia koa* (Meyrick 1928, Swezey 1954), and are parasitized by the ichneumonid wasp *Pristomerus hawaiiensis* Perkins (Zimmerman 1978). Adults are readily attracted to ultraviolet lights, typically several hours after sunset, in dense or mature *Acacia koa* forests with senescing branches.

DISTRIBUTION: Endemic to Hawaiian Islands: Maui, Oahu, Kauai. The type specimen is the only known specimen from Maui. This species may persist at low densities on Maui, but probably would be in competition with *C. acaciavora*. This species is relatively common in dense mature *Acacia koa* stands on Kauai.

REMARKS: The forewing patterns in this species are somewhat variable requiring genitalia preparations for conclusive identification. Zimmerman (1978:599) suggested the approximation of the signa in the corpus bursae is a diagnostic characteristic for this species. However, closer examination of the holotype slide (BM 2052) reveals that the signa are on opposite sides of the corpus bursae giving the appearance of proximity. Zimmerman's own slide preparation (Z-XII-62-5) appears to be misplaced (at BPBM) and therefore unavailable for examination, but illustrates the same illusion (Zimmerman, 1978, Figure 391). The male genitalia slide figured by Zimmerman (1978, Figure 385) is also misplaced (at BPBM), but shows a bilobed uncus margin not seen in other specimens of this species.

Cydia rufipennis (Butler 1881)
(Figures 42, 43, 61, 80)

Phoxopteris rufipennis Butler 1881:395

Adenoneura rufipennis. – Walsingham 1907:680; Plate X. Figure 22 (moth), Swezey 1936:198, 1954:4

Cydia rufipennis. – Zimmerman 1978:609; Figures 375 (wing venation), 382 (moth), 388 (♂ genitalia), 397 (♀ genitalia), Plate 2:8 (moth). Stein 1983a:318, 1983b:300. Miller 1990:136

DIAGNOSIS: *Cydia rufipennis*, from Oahu and Kauai and its sister, *C. montana*, from Hawaii and Maui, are the two smallest species of Hawaiian *Cydia*. They appear to be in the early stages of diversification and may represent an evolutionary grade rather than two separate clades. The size and reddish hue of *C. rufipennis* is unique among Hawaiian *Cydia*.

DESCRIPTION: (exp. 6-10 mm, n=26) **Head:** Head testaceous, labial palpi buff with testaceous flecks and testaceous third segment, antennae banded ferruginous and buff. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally ferruginous scales with buff tips, mixed with occasional testaceous scales, particularly in scutellar area, tegulae same. Without dorsal tuft of scales. Ventrally whitish-buff. Fore- and middle-legs ferruginous scales with buff tips, middle-tibia testaceous, middle and hind-tarsi banded fuscous and buff. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figures 42-43) Slightly dilated distally, costa gently arched, apex obtuse, termen nearly straight. Ground color fuscous-brown. Costal strigulae distinct interspersed with testaceous streaks, directed distally towards termen, striae indistinct in basal half creating mottled appearance. No triangular costal patch. Testaceous oblique medial fascia extending from end of cell towards basal dorsum, bordered distally by small whitish discal patch. Testaceous streak extending from medial fascia below discal patch, connecting with vague testaceous pretornal blotch. Ocellar patch light ferruginous with 2-3 ocellar spots, bordered distally by a plumbaginous crescent and anteriorly by oblique testaceous streak ending in the central termen. Testaceous subtriangular apical patch extends along termen as a thin band. Fringe ferruginous. Ventrally costal strigulae apparent along costal margin, otherwise uniform dark brown. **Hindwings:** Dorsally and ventrally dark brown, fringe light brown-buff. Males with glandular ventral pouch along path of CuP opening dorsally and enclosing elongate modified pecten scales. Vein A3 in males displaced towards anal margin. Males without anal roll or androconial scales (thecae) along vein A3. **Abdomen:** Uniformly dark brown, caudal area sometimes ringed by light brown-buff scales. **Male genitalia:** (Figure 61) Tegumen simple with small central point along caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa gently concave, ventral edge with somewhat shallow invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with short setae along ventral and distal margin sparsely intersperse with longer setae. Aedeagus simple, curved, without cornuti, tip with a slight dorsal notch. **Female genitalia:** (Figure 80) Lamellae postvaginalis longer than wide, constricted centrally and flared into two lobes on the caudal end. Antrum narrow elongate cylindrical with no obvious sclerotization, not extending past junction with ductus bursae. Corpus bursae with diverticulum, two falcate signa. Ductus bursae shorter

than width of corpus bursae, with broad junction to corpus bursae, but otherwise narrow as antrum.

TYPE MATERIAL: *Phoxopterus rufipennis* – ♀ holotype (BMNH): Hawaiian Islands, 81.7 109 [Blackburn code on other specimens indicating “the mountains of Oahu” Zimmerman (1978: 609)]. ♂ cenotype (BMNH): Oahu, Waianae Mtns., 2000 ft, IV.1892, RCL Perkins; Walsingham specimen 25094. ♀ cenotype (BMNH): Oahu, Waianae Mtns., 2000 ft, IV.1892, RCL Perkins; Walsingham specimen 25097.

ADDITIONAL MATERIAL: 1♂ (BMNH): Hawaiian Islands, Kauai, Lihue, 4000 ft, VII.1896, RCL Perkins; Walsingham specimen 28032; genitalia slide BM 2058. 1♀ (BMNH): Hawaiian Islands, Kauai, Kaholuamano, 4000 ft, IV.1895, RCL Perkins; Walsingham specimen 27734. 1♂ (BMNH): Hawaiian Islands, Oahu, Halemano, c. 2000 ft, XII.1892; RCL Perkins; Walsingham specimen 25822. 1♀, 1? (BMNH): Oahu, Waianae Mtns., 2000 ft, IV.1892, RCL Perkins; Walsingham specimens 25096, 25100 respectively. 2 (BMNH): Hawaiian Islands, 1899, Blackburn. 3♀ (BPBM): Oahu, Waianae Mtns., 2000 ft, IV.1892, RCL Perkins; Walsingham specimen 25095, 25098, 25099; (abdomen missing from 25095 & 25098). 1♂ (BPBM): Oahu, Honolulu, 21.iii.1917, JC Bridwell. 1♀ (BPBM): Oahu, Tantalus, OH Swezey; (right FW missing). 1♂, 1♀ (BPBM): Oahu, Kahili, ex: *Acacia koa* seeds, OH Swezey; (♀ abdomen missing). 1♂ (EMEC): Hawaiian Islands, Kauai, Kokee State Park, Awaawapuhi trail, 1101 m., at UV light, 23.vi.2005, PT Oboyski. 3♀ (EMEC): Hawaiian Islands, Kauai, Kokee State Park, 4WD road off Canyon road, 889 m., at UV light, 25.vi.2005, PT Oboyski. 1♂ (UHMA): HI: Kauai Island, Na Pali - Kona F.R., Mohihi Rd @ Waiakoali Stream, 3400 ft., V-17-05, D. Rubinoff et al. 1♂ (UHMA): HI: Oahu, Koolau Mtns., Wiliwilinui Trail, 6 June 2008, UV light trap, J. Eiben & W. Haines.

BIOLOGY: Larvae of *C. rufipennis* feed on *Acacia koa* within developing seeds, flowers, and flower buds (Swezey 1936, 1954, Zimmerman 1978, Stein 1983a, b). Bridwell (1919) reports that last instar larvae emerge from seedpods to pupate elsewhere. The adults I have collected on Kauai at ultraviolet lights were only amid flowering *Acacia koa* trees.

DISTRIBUTION: Endemic to Hawaiian Islands: Kauai (particularly around Kokee State Park) and Oahu – widely distributed but only locally abundant where *Acacia koa* grows.

REMARKS: *Cydia rufipennis* from the older islands (Kauai and Oahu) is sister to *C. montana* on the younger islands (Maui and Hawaii), and appears to occupy a similar niche.

Cydia montana (Walsingham 1907)
(Figures 44, 45, 46, 62, 81)

Adenoneura montanum Walsingham 1907:679; Plate X. Figure 21 (moth). Swezey & Williams 1932:187, Swezey 1936:198, 1954:204

Cydia montana. – Zimmerman 1978:595; Figures 373 (wing venation), 381 (moth), 385 (♂ genitalia), 393 (♀ genitalia). Stein 1983a:318

DIAGNOSIS: *Cydia montana*, from Hawaii and Maui, and its sister *C. rufipennis*, from Oahu and Kauai, are the two smallest species of Hawaiian *Cydia* (exp. 8-11 mm and 6-10 mm, respectively). They appear to be in the early stages of diversification and may represent an evolutionary grade rather than two separate clades. The light brown mottled coloring of *C. montana* resembles a diminutive *C. storeella* or *C. plicata*, compared to the reddish hue of *C. rufipennis*. However, larger species, such as *C. plicata*, can have diminutive forms under poor food resource conditions (Oboyski, personal observation), as has been noted for other *Cydia* species (Miller 1990). *Cydia montana* is easily distinguished from *C. rufipennis* by the distribution and red hue of the latter, and from *C. storeella* and *C. plicata* by the shape of the female antrum.

DESCRIPTION: (exp. 8-11 mm, n=25) Colored scales on head, body, and legs mostly shades of ferruginous-brown with whitish tips. **Head:** Antennae, head and labial palpi light brown-cinereous. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally light ferruginous-brown, tegulae same. Without dorsal tuft of scales. Ventrally lighter ferruginous-brown. . Legs light brown-ochreous, tarsomeres banded dark brown-cinereous at bases, lighter distally. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figure 44-46) Slightly dilated distally, costa gently arched, apex obtuse, termen nearly straight. Ground color light brown-ochreous. Costal strigulae distinct, directed distally towards termen, striae indistinct in basal half creating mottled appearance. No triangular costal patch. Oblique medial fascia and pretornal blotch indistinct. Whitish discal patch faint. Ocellar patch light brown-ochreous with three ocellar spots, bordered distally by a light bronze crescent. Apex often with a light ferruginous-brown patch extending into fringe. Continuation of ferruginous-brown stria to central termen extends into fringe. Fringe otherwise ferruginous-white. Ventrally dark brown-cinereous, lighter along anal margin. **Hindwings:** Dorsally cinereous-brown basally, ferruginous-brown distally. Males with glandular ventral pouch along path of CuP opening dorsally and enclosing elongate modified pecten scales. Vein A3 in males displaced towards anal margin. Males without anal roll or androconial scales (thecae) along vein A3. **Abdomen:** Uniformly dark brown-cinereous dorsally, lighter ventrally. **Male genitalia:** (Figure 62) Tegumen simple with somewhat bilobed crista along caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa gently concave, ventral edge with somewhat shallow invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with relatively sparse short setae along ventral and distal margin intersperse with longer setae relative to other Hawaiian *Cydia*. Aedeagus simple, curved, without cornuti, and somewhat bilobed at tip. **Female genitalia:** (Figure 81) Lamellae postvaginalis slightly longer than wide, hardly constricted centrally. Antrum narrow elongate cylindrical with no obvious sclerotization, not extending past junction with ductus bursae. Corpus bursae with diverticulum, two long falcate signa. Ductus

bursae much shorter than width of corpus bursae, with broad junction to corpus bursae, but otherwise narrow as antrum.

TYPE MATERIAL: *Adenoneura montanum* – ♂ holotype (BMNH): Hawaii, Mt Kilauea, vii.1895, RCL Perkins; Walsingham specimen 27483; genitalia slide BM 2055. ♀ holotype (BMNH): Hawaii, Kona, 4000 feet, 10.viii.1892, RCL Perkins; Walsingham specimen 25275. 2♂, 2♀ paratypes (BMNH): Hawaii, Mt Kilauea, vii.1895, RCL Perkins; Walsingham specimens 27423, 27396, 27398, 27405, respectively; ♂ wing slide BM 7531; ♀ genitalia slide BM 2056. ♀ paratype (BMNH): Hawaii, Kona, 4000 ft, 8.ix.1892, RCL Perkins; Walsingham specimen 25561. 3♂ paratypes (BPBM): Hawaii, Mt Kilauea, vii.1895 (2 specimens), viii.1896 (1 specimen), RCL Perkins, Walsingham specimens 27412, 27403, 28083, respectively. 1♀ paratype (mistakenly labeled by Walsingham as ♂) (BPBM): Hawaii, Mt Kilauea, viii.1895, RCL Perkins, Walsingham specimens 27464. ♀ allotype (BMNH?): Hawaii, Kona, 4000 feet; (abdomen lost).

ADDITIONAL MATERIAL: 1 (HDOA): Hawaii, Nauhi gulch, 5000-6000 ft, koa?, 30.ix.1931, OH Swezey & FX Williams; determined by Swezey. 3 (HDOA): Hawaii, Kilauea, HAVO, ex: light trap, ix.1946, CJ Davis; determined by Swezey. 1♀ (EMEC): Hawaiian Islands, Hawaii, Mauna Kea, Hakalau N.W. R., Nauhi cabin, 1640 m., at mercury vapor light, 8.viii.2003, PT Oboyski. 1♀ (EMEC): Hawaiian Islands, Hawaii, Kilauea, HAVO N.P., residential quarters 20, 1150 m., at mercury vapor light, 5.vii.2004, PT Oboyski. 4♂,6♀ (EMEC): Hawaiian Islands, Hawaii, Mauna Loa, HAVO N.P., Mauna Loa strip road, 1885 m., at UV light, 2.viii.2005, PT Oboyski. 3♂ (EMEC): Hawaiian Islands, Hawaii, Kilauea, HAVO N.P., crater rim trail nr. military camp, 1200 m., at UV light, 4.viii.2005, PT Oboyski. 3♀ (EMEC): Hawaiian Islands, Hawaii, Hualalai, top of Kaloko Drive, 1485 m., at UV light, 5.viii.2005, PT Oboyski. 2♀ (EMEC): Hawaiian Islands, Hawaii, Mauna Loa, Kipuka Puu Huluhulu (west side), 2010 m., at UV light, 10.viii.2005, K. Roggeveen. 5♂,2♀ (EMEC): Hawaiian Islands, Hawaii, Mauna Loa, Saddle Road Kipuka mp 21, 1695 m., at UV light, 12.viii.2005, K. Roggeveen. 1♂,1♀ (EMEC): Hawaiian Islands, Hawaii, Mauna Loa, HAVO N.P., Mauna Loa strip road, 1940 m., at UV light, 17.viii.2005, K Roggeveen.

BIOLOGY: Zimmerman (1978) and Stein (1983a), following Swezey (1936, 1954), lists *Sophora chrysophylla* seeds as the host of *C. montana*. I have not been able to locate the specimens Swezey claims to have reared from *Sophora* and suspect that he was mistaken. In fact, one specimen collected by O.H. Swezey and F.X. Williams from Nauhi Gulch, Hawaii Island includes “koa?” on the collection label. Although I have not reared this species, I have observed large numbers of *C. montana* adults at ultraviolet lights located amid mature *Acacia koa* trees, but never among *Sophora* trees. I have also swept adults of this species from the canopy of *Acacia koa* at 1715 m elevation at Keauhou Ranch, Mauna Loa. Extensive efforts to rear larvae from *Acacia koa* seedpods and rust galls have produced mostly *Cryptophlebia illepada* (Butler) and few *Cydia walsinghamii* (Butler), but no *C. montana*. Less effort was spent rearing larvae from *Acacia* flowers, a known substrate for *C. rufipennis*, the sister to *C. montana*. Based on this strong circumstantial evidence, I suspect *C. montana* feeds on *Acacia koa*, particularly on flowers.

DISTRIBUTION: Endemic to Hawaiian Islands: Hawaii – distributed around the island along the “koa belt” (600-1200 m). One specimen collected from 1290 m on Haleakala, Maui.

REMARKS: Although previously thought to feed on *Sophora*, this species is associated with *Acacia koa* and is the Hawaii Island (and Maui) sister-species of *C. rufipennis* found on the older islands (Oboyski Chapter 3). The one specimen I collected on Maui has a reddish hue similar to *C. rufipennis* but the mitochondrial DNA haplotype of *C. montana* (Oboyski Chapter 3), and appears to represent an intermediate form. Specimens misidentified as *C. montana* by Brenner et al. (2002) are correctly referred to *C. plicata*.

Cydia hawaiiensis n. sp.

(Figures 47, 65, 83)

DIAGNOSIS: *Cydia hawaiiensis* is easily confused with *C. walsinghamii* based on size, markings, and habitat, and probably is a close relative. Males of *C. hawaiiensis* have a very shallow pouch on the hindwing relative to other Hawaiian *Cydia*. Males of *C. walsinghamii* lack a hindwing pouch. The genitalia of *C. hawaiiensis* females are nearly identical to *C. walsinghamii*.

DESCRIPTION: (exp. 18-22 mm, n=11) **Head:** Antennae light brown-ferruginous. Head and labial palpi pale ochreous-buff. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally pale ochreous-buff mottled with fuscous and ferruginous, tegulae with ferruginous bases grading to ochreous caudally. Without dorsal tuft of scales. Venter whitish-buff. Forelegs fuliginous, fore- and midtibia and tarsi banded fuliginous and ochreous. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figure 47) Slightly dilated distally, costa gently arched, apex obtuse, termen nearly straight. Ground color buff. Costal strigulae distinct, directed distally towards termen. Striae in basal area indistinct, mottled ochreous, fuscous, and ferruginous. Triangular costal patch absent. Oblique medial fascia interrupted, represented by dark brown-fuscous patch towards end of cell and oblique patch from fold to dorsum, bordered distally by small white discal patch and whitish oblique band. Costal strigulae in distal half of costa sometimes suffused with dark brown-fuscous patch connecting to dark patch at end of cell. Pretornal blotch reaching to dark patch at end of cell, bordered by dark brown-fuscous with ferruginous center. Ocellar patch light brown-ochreous, ocellar spots indistinct, bordered distally by thin plumbaginous crescent. Dark brown apical patch suffused over much of apical area extending into fringe. Fringe at mid-termen and tornus light brown-ochreous. Ventrally, costal strigulae apparent along costa, otherwise uniformly fuscous-brown. **Hindwings:** Uniformly brown dorsally and ventrally. Males with a very shallow glandular ventral pouch along path of CuP opening dorsally and enclosing elongate modified pecten scales. Vein A3 in males displaced towards the anal margin. Males without an anal roll or androconial scales (thecae) along vein A3. Abdomen: Uniformly dark brown. **Male genitalia:** (Figure 65) Tegumen simple, with broadly rounded caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa slightly concave, ventral margin with deep invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with dense short and long setae along ventral and distal margin. Aedeagus simple, curved, without cornuti, tip split by two dorsal ridges. **Female genitalia:** (Figure 83) Lamellae postvaginalis slightly longer than wide, with slight central constriction.

Antrum elongate funnel-shaped with small sclerotized line near junction with ductus bursae. Corpus bursae with diverticulum and two falcate signa. Ductus bursae shorter than width of corpus bursae, broadly joined with corpus bursae, quickly tapering to antrum.

TYPE MATERIAL: *Cydia hawaiiensis* – ♂ holotype (BPBM): Hawaiian Islands, Hawaii, Mauna Loa, HAVO N.P., Mauna Loa trail, *Acacia koa* forest, 2035 m., at UV light, 5.vii.2004, PT Oboyski; Oboyski specimen PTO-279.8, genitalia slide pto-s304. ♀ allotype (BPBM): Hawaiian Islands, Hawaii, Mauna Loa, HAVO N.P., Mauna Loa strip road top, *Acacia koa* forest, 2180 m., at UV light, 17.vii.2004, PT Oboyski; Oboyski specimen PTO-322.2, genitalia slide pto-s042. 1♂ paratype (EMEC): Hawaiian Islands, Hawaii, Mauna Loa, Papaloa (upslope of Kona), *Acacia koa* forest, 1560 m., at UV light, 9.vii.2003, PT Oboyski; Oboyski specimen PTO-218.2, genitalia slide pto-s301. 10♂ paratypes (EMEC): Hawaiian Islands, Hawaii, Kilauea, HAVO N.P., Kipuka Puauulu, *Acacia koa* forest, 1195 m., at UV light, 15.i.2004, PT Oboyski. Oboyski specimens PTO-238.1-10, genitalia slides pto-s302, s303.

ADDITIONAL MATERIAL: This species is known only from the type material although some female specimens labeled as *Cydia walsinghamii* in existing collections may be *C. hawaiiensis*.

BIOLOGY: This species is known only from adults collected at ultraviolet lights in *Acacia koa* forest. Probably larvae feed on *Acacia koa*.

DISTRIBUTION: Endemic to Hawaiian Islands: Hawaii.

REMARKS: Adults are attracted to ultraviolet lights.

Cydia acaciavora n. sp.
(Figures 48, 66, 84)

DIAGNOSIS: *Cydia acaciavora* is one of four *Cydia* species in Hawaii (*C. acaciavora*, *C. anomalosa*, *C. crassicornis*, *C. walsinghamii*) that lack the pouch in the male hindwing. Males of *C. acaciavora* can be separated from these other three species by the genitalia, particularly by the long slender shape of the aedeagus. Females of *C. crassicornis* are not known and *C. anomalosa* differ in the shape of the antrum. Females of *C. acaciavora* and *C. walsinghamii* are difficult to distinguish, although the antrum of *C. acaciavora* is more narrow than that of *C. walsinghamii*.

DESCRIPTION: (exp. 13-17 mm, n=4) **Head:** Antennae light brown, head and labial palpi pale ochreous, palpi becoming fuscous distally and ventrally. Palpi slightly upcurved, third segment projecting beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally dark brown-fuscous, tegulae pale ochreous. Without dorsal tuft of scales. Venter whitish-buff. Legs pale cinereous, fore- and midtibia and tarsi banded fuscous and pale ochreous. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figure 48) Slightly dilated distally, costa gently arched, apex obtuse, termen nearly straight. Ground color whitish-buff. Costal strigulae distinct, directly distally towards termen. No triangular costal patch. Basally mottled pale fuscous above cell, cell and below buff mottled with thin fuscous streaks. Oblique medial fascia of light brown vague, with darker distal edge, bordered distally by whitish band and whitish discal patch. Pretornal blotch of dark brown-ferruginous broad and short. Subtriangular

apical patch dark brown-ferruginous extends into fringe. Ocellar patch whitish with 2-3 indistinct ocellar spots, bordered distally by silvery-white crescent. Continuation of brown-ferruginous stria to central termen extends into fringe. Fringe otherwise silvery-buff. Ventrally brown-ferruginous with costal strigulae somewhat apparent. **Hindwings:** Uniformly brown-ferruginous. Males without a glandular ventral pouch, anal roll, or androconial scales (thecae) along vein A3. **Abdomen:** Uniformly brown-ferruginous. **Male genitalia:** (Figure 66) Tegumen simple, with broadly rounded caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa gently concave, ventral margin with deep invagination. Saccus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with dense short setae along ventral and distal margin interspersed with longer setae. Aedeagus simple, curved, long and slender with wider base, without cornuti, flared spatulate tip with dorsal excavation for one-fifth length of aedeagus. **Female genitalia:** (Figure 84) Lamellae postvaginalis longer than wide and relatively narrowed by central constriction. Antrum relatively short and narrow with sclerotized ring at junction with ductus bursae. Corpus bursae with diverticulum and two falcate signa. Ductus bursae short, joined broadly to corpus bursae, quickly tapering to antrum.

TYPE MATERIAL: *Cydia acaciavora* – ♂ holotype (BPBM): MAUI, Haleakala N.P., Kipahulu Vy., Delta Camp. UV light, 15 Jan 2004, D. Rubinoff, W. Haines; Oboyski specimen PTO-343.5, genitalia slide pto-s316. 2♀ paratypes (EMEC): MAUI, Haleakala N.P., Kipahulu Vy., Delta Camp. UV light, 15 Jan 2004, D. Rubinoff, W. Haines; Oboyski specimens PTO-343.1, genitalia slide pto-s315, PTO-343.2. 1♂ paratype (EMEC): Hawaiian Islands, Maui, Olinda, Makawao F.R., Kula pipeline trail, 1280 m., reared from dead *Acacia koa* branch, 27.vii.2002, PT Oboyski; Oboyski specimen PTO-95.20. 1♂ paratype (EMEC): Hawaiian Islands, Hawaii, Kohala, Kawaihae Uka, *Acacia koa* sanctuary, 975 m., sweeping *Acacia koa*, 5.vii.2003, PT Oboyski; Oboyski specimen PTO-149.1, genitalia slide pto-s307.

BIOLOGY: Larvae feed under bark of senescing branches and in *Uromyces koae* rust galls on *Acacia koa*. Although *Pristomerus hawaiiensis* emerged from rearing substrates, it was not possible to determine whether they were associated with *Cydia acaciavora*, *Cydia walsinghamii*, or *Cryptophlebia illepidata* larvae, all of which produced adults from the same host material. Adults are attracted to ultraviolet lights.

DISTRIBUTION: Endemic to Hawaiian Islands: Maui – Haleakala. Hawaii – One specimen collected from Kawaihae Uka, *Koaia* sanctuary.

REMARKS: This species may be more widespread than realized, but dismissed as a form of *C. walsinghamii*. Mitochondrial DNA evidence corroborates these are two separate species (Oboyski Chapter 3).

Cydia walsinghamii (Butler 1882)
(Figures 49, 67, 85)

Proteopteryx walsinghamii Butler 1882:43

Enarmonia walsinghami. – Walsingham 1907:684, 736; Plate XI. Figure 1 (moth). Swezey 1954:5

Cydia walsinghamii. – Zimmerman 1978:610; figures 375 (wing venation), 383 (moth), 388 (♂ and ♀ genitalia)

DIAGNOSIS: *Cydia walsinghamii* is one of four *Cydia* species in Hawaii (*C. acaciavora*, *C. anomalosa*, *C. crassicornis*, *C. walsinghamii*) that lack the pouch in the male hindwing. This species can be distinguished by the shape of the antrum in females and the tip of the aedeagus in males.

DESCRIPTION: (exp. 10-22 mm, n=37) **Head:** Antennae, head, and labial palpi light brown-cinereous. Palpi slightly upcurved, third segment projecting beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally light brown-cinereous gradually becoming darker towards scutellum, tegulae same. Without dorsal tuft of scales. Venter light brown-buff. Legs light brown cinereous, hind tarsi paler. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figure 49) Slightly dilated distally, costa gently arched, apex obtuse, termen slightly sinuous. Ground color whitish-buff. Costal strigulae distinct, directed distally towards termen. The following patterns variable in color and degree of suffusion. Triangular costal patch of light brown-ferruginous to dark brown sometimes present, suffusing through some of the following patterns. Basal area light brown cinereous mottle with streaks of dark brown-ferruginous. Oblique medial fascia ferruginous with dark brown-ferruginous streaks from end of cell towards basal dorsum, bordered distally by band of whitish-buff and whitish discal patch. Conspicuous pretornal blotch of ferruginous to dark brown-ferruginous nearly reaching to discal patch. Ocellar patch whitish-buff with three distinct ocellar spots, bordered distally by silvery-white crescent. Subtriangular apical patch ferruginous extending into fringe. Continuation of brown-ferruginous stria to central termen extends into fringe. Fringe otherwise pale cinereous. Ventrally brown-cinereous. **Hindwings:** Uniformly light brown-cinereous to fuscous. Males without a glandular ventral pouch, anal roll, or androconial scales (thecae) along vein A3. **Abdomen:** brown-cinereous. **Male genitalia:** (Figure 67) Tegumen simple, crista of broadly rounded caudal ridge with central point, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa slightly concave, ventral margin with deep invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with dense short and long setae along ventral and distal margin. Aedeagus simple, slightly curved, stout near base and middle tapering quickly to end with shallow dorsal excavation at tip, without cornuti. **Female genitalia:** (Figure 85) Lamellae postvaginalis slightly longer than wide, constricted centrally. Antrum funnel-shaped with one thin line of sclerotization on either side and a sclerotized ring at junction with ductus bursae. Corpus bursae with diverticulum and two falcate signa. Ductus bursae shorter than width of corpus bursae, broadly joined with corpus bursae, quickly tapering to antrum.

TYPE MATERIAL: *Proteopteryx walsinghamii* – ♀ holotype (BMNH): Hawaiian Islands, 89.2; 161 [Blackburn collecting code indicating “in the neighborhood of Honolulu” (Zimmerman, 1978:610)]; genitalia slide BM 3756. ♂ cenotype (BMNH): Mt Kilauea, Hawaii, VIII.1896, RCL Perkins; Walsingham specimen 28075; genitalia slide BM 1892. ♀ cenotype (BMNH): Kona, 4000 ft, Hawaii, 12.VIII.1892, RCL Perkins; Walsingham specimen 25765; genitalia slide BM 1893.

ADDITIONAL MATERIAL: 1♀ (BPBM): Hawaii, Hilo, 2000 ft, xii.1895, RCL Perkins; Walsingham specimen 28599. 1♀ (BPBM): Hawaii, Hualalai (Kona), 5000 ft, 5.viii.1892, RCL Perkins; Walsingham specimen 25830; (abdomen missing). 1♂ (BPBM): Hawaii, Kona, 4000 ft, 10.viii.1892, RCL Perkins; Walsingham specimen 25747. 4♂ (EMEC): Hawaiian Is: Hawaii, M. Loa, Kipuka Puu Huluhulu, *A. koa* forest – UV light, 2005 m., 12.viii.2005, K. Roggeveen. 4♂ (EMEC): Hawaiian Is: Hawaii, M.Loia, HAVO N.P., Mauna Loa strip rd, *A.koa* forest – UV light, 1885 m., 2.viii.2005, PT Oboyski. 5♂ (EMEC): Hawaiian Is: Hawaii, Hualalai, top of Kaloko Drive., *A.koa* forest – UV light, 1485 m., 5.viii.2005, PT Oboyski. 7♂ (EMEC): Hawaiian Is: Maui, Haleakala, upper Makawao F.R., *A.koa* forest – UV light, 1291 m., 3.vii.2005, PT Oboyski. 6♂,5♀ (EMEC): Hawaiian Is: Kauai, Kohee, Awaawapuhi trail, ~1.5 mi., *A.koa* forest – UV light, 1027 m., 21.vi.2005, PT Oboyski.

BIOLOGY: Larvae of *Cydia walsinghamii* feed on *Acacia koa* and *A. koaia* in dead twigs, living tips of twigs, and galls caused by *Uromyces koae* rust fungus (Swezey 1954), as well as *Acacia koa* seeds (Zimmerman 1978). Larvae compete for these resources with *Cryptophlebia illepada*, *Cydia rufipennis*, *C. montana*, and *C. conspicua*, although some resource partitioning likely takes place among species. *C. walsinghamii* larvae are parasitized by the ichneumonid *Pristomerus hawaiiensis* and probably other parasitoids that attack *Cydia* larvae in Hawaii. Adults are often abundant at lights in *Acacia* habitat.

DISTRIBUTION: Endemic to Hawaiian Islands: Common and abundant where mature *Acacia koa* trees can be found on Hawaii, Maui, Oahu, and Kauai islands. It probably occurred on Molokai and Lanai, but naturally growing *A. koa* no longer occurs on these islands. However, I reared one specimen from a dead twig of *Acacia koaia* from a senescing natural population of this tree within a protective enclosure fence in the Kamiloloa section of the Molokai Forest Reserve, Molokai.

REMARKS: This is a widespread and extremely variable species, both in size and color. Zimmerman (1978) suspected *C. walsinghamii* to be a species complex, but did not attempt to qualify the different forms. The collection of Dr. Klaus Sattler at the BMNH contains long series of morphologically variable specimens from Hawaii, Oahu, and Kauai that may represent new species. However, these specimens were not available for dissection and genitalia analysis. More than one species may exist within what is currently called *walsinghamii*, but this is obscured by the extreme variation within this species.

Cydia crassicornis (Walsingham 1907)
(Figure 68)

Enarmonia crassicornis Walsingham 1907:685; Plate XI. Figure 2 (moth)

Cydia crassicornis. – Zimmerman 1978:585; Figures 373 (wing venation), 378 (moth), 384 (male genitalia)

DIAGNOSIS: *Cydia crassicornis* is one of four *Cydia* species in Hawaii (*C. acaciavora*, *C. anomalosa*, *C. crassicornis*, *C. walsinghamii*) that lack the pouch in the male hindwing. Although possibly a form of the widespread and polymorphic *walsinghamii*, the caudal ridge of the tegumen lacks the minute crista (Figure 68) found on *walsinghamii* (Figure 67). The tip of the aedeagus of male *C. crassicornis* (Figure 68) is not excavated as in *C. acaciavora* (Figure 66). Males of *C. crassicornis* also lack the hindwing anal roll and thecae along vein A3 as in *anomalosa*.

DESCRIPTION: (exp. 14 mm, n=2) **Head:** Antennae light brown-cinereous, labial palpi buff white dorsally, light brown-cinereous laterally. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Head buff white, post ocular scales light brown-cinereous, some scales with dark tips. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally and ventrally buff white, mid-dorsum light brownish white, tegulae light ferruginous anteriorly, becoming lighter posteriorly. Without dorsal tuft of scales. Legs (very worn) light brown-cinereous with cinereous-white spurs and tarsi, midtibia darker with light-tipped scales. No discernable sex scales (e.g. hair pencils). **Forewings:** Slightly dilated distally, costa gently arched, apex barely obtuse, termen straight or slightly convex. Ground color buff white. Costal strigulae distinct, directed distally towards termen. Triangular costal patch mottled fuscous-ferruginous suffused from base to 2/3 length of costa, posteriorly overlapping the fold. Dorsal area light ochreous-white contiguous with white patch at distal end of cell, very light brown-cinereous suggestion of a pretornal patch. Ocellar patch with two or three indistinct ocellar spots bordered distally by silvery-bronze crescent. Apex with a light fuscous patch extending into fringe. Continuation of fuscous-ferruginous stria to central termen extends into fringe. Ventrally brown-fuscous. **Hindwings:** Uniformly light brown-cinereous dorsal and ventral, fringe whitish-gray. Males without glandular pouch, cubital pecten not modified. Vein A3 in males similar to females of other species, not displaced towards anal margin as in species with glandular pouch. Males without anal role or androconial scales (thecae) along vein A3. Abdomen: Uniformly brown-cinereous. **Male genitalia:** (Figure 68) Tegumen simple, broadly rounded along caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa slightly concave, ventral margin with deep invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with dense short setae along ventral and distal margin, interspersed with fewer long setae. Aedeagus simple, slightly curved, without cornuti, tip slightly excavated dorsally with short dorsal projection. **Female genitalia:** (females not known).

TYPE MATERIAL: *Enarmonia crassicornis* Walsingham – ♂ holotype (BMNH): Hawaii, Kona, above 4000 ft., ix.1892, RCL Perkins; Walsingham collection 25212; genitalia slide BM 1880. ♂ paratype (BMNH): Hawaii, Kona, above 4000 ft., ix.1892, RCL Perkins; Walsingham collection 25215.

ADDITIONAL MATERIAL: This species is known only from two males collected by R.C.L. Perkins in 1892.

BIOLOGY: Perkins' field notes for September 1892 indicate he "collected ... by sifting dead leaves at the foot of a big koa tree (4000 ft.)," and later that month in the "rain belt" at lower elevations above Kona (Evenhuis 2007). Perkins also collected the koa-feeding *C. montana* at 4000 ft. above Kona in August and September 1892. Although it is likely *C. crassicornis* was associated with *Acacia koa*, without further information, it is not possible to assign the proper host plant, habitat, larval biology, predators, or parasitoids for this species.

DISTRIBUTION: Endemic to Hawaiian Islands: Hawaii Island – known only from two specimens collected above Kona above 1200 m.

REMARKS: If not extinct, this species is very rare. The wing venation and genitalia confirm the placement of this species within *Cydia*, though it is unclear without more material whether it represents a separate lineage in Hawaii.

***Cydia anomalosa* n. sp.**
(Figures 19, 21, 23, 50, 51, 69, 86)

DIAGNOSIS: Males of *C. anomalosa* are the only *Cydia* in Hawaii with an anal roll on the hindwing, a characteristic common among non-Hawaiian *Cydia* species. Males also lack the hindwing pouch found on many other Hawaiian *Cydia*. Females can be distinguished from the similar-looking *C. walsinghamii* by the shape of the female antrum.

DESCRIPTION: (exp. 10-13 mm, n=11) **Head:** Antennae, head, and labial palpi light brown-cinereous. Palpi slightly upcurved, third segment projecting forward beyond vestiture of second segment. Ocelli and chaetosemata conspicuous. **Thorax:** Dorsally and ventrally light brown-cinereous, tegulae same. Without dorsal tuft of scales. Legs light brown-cinereous. No discernable sex scales (e.g. hair pencils). **Forewings:** (Figures 50-51) Slightly dilated distally, costa gently arched, apex obtuse, termen somewhat sinuate. Ground color whitish-buff. Costal strigulae distinct, directed distally towards termen, sometimes obscured by triangular costal patch of brown-cinereous from base to two-thirds length of costa, posteriorly overlapping the fold. Oblique medial fascia of dark-brown cinereous extending from end of cell towards basal dorsum. Occasionally anal area below fold to pretornal blotch suffused whitish-buff. Pretornal blotch indistinct area of light brown to coppery-brown. Whitish discal patch often indistinct. Ocellar area olivaceous to light brown with three ocellar spots, bordered distally by silvery crescent. Subtriangular apical patch of light brown to coppery brown sometimes extending into fringe. Continuation of ferruginous-brown stria to central termen sometime extending into fringe. Termen with thin band of dark brown-ferruginous from apex through tornus, sometime interrupted. Fringe otherwise light brown-cinereous. Ventrally brown-fuscous. **Hindwings:** Ventrally brown-fuscous, lighter along costal area. Males with anal roll enclosing whitish scales within (Figure 19) and phylliform scales along anal margin (Figures 21, 23). Males without glandular ventral pouch or modified pecten scales. **Abdomen:** brown-cinereous. **Male genitalia:** (Figure 69) Tegumen simple, with broadly rounded caudal ridge, lacking gnathos, socii, and uncus. Valvae with excavation in basal third, costa slightly concave, ventral margin with deep


invagination. Sacculus slightly sinuous, with slight concavity near base. Cucullus elongate rounded with short and long setae intersperse along ventral and distal margin. Aedeagus simple, curved, relatively short and stout, tip mostly flat with a dorsal “hood”. **Female genitalia:** (Figure 86) Lamellae postvaginalis nearly as wide as long, constricted centrally. Antrum simple cylindrical connecting directly with ductus bursae. Corpus bursae with diverticulum and two falcate signa. Ductus bursae shorter than width of corpus bursae, broadly connecting with corpus bursae and tapering only slightly to antrum.

TYPE MATERIAL: *Cydia anomalosa* – ♂ holotype (BPBM): Hawaiian Islands, Kauai, Mt. Kahili State Forest Reserve, 354 m., at UV light, 24.vi.2005, PT Oboyski; Oboyski specimen PTO-513.3. 7♂,2♀ paratypes (EMEC): Hawaiian Islands, Kauai, Mt. Kahili State Forest Reserve, 354 m., at UV light, 24.vi.2005, PT Oboyski; Oboyski specimens PTO-513.1,2,4,5,6,7,8,9,14, genitalia slides pto-s163 (♂ PTO-513.1), pto-s164 (♀ PTO-513.2), pto-s169 (♀ PTO-513.5). 1♂ paratype (EMEC): Hawaiian Islands, Maui, Makawao F.R., Kula pipeline trail, 1280 m., reared from *Acacia koa* pods, 27.vii.2002, PT Oboyski; Oboyski specimen PTO-096.5. 1♂ paratype (EMEC): Hawaiian Islands, Maui, Haleakala N.P., Kaupo Gap, 1680 m., reared from *Acacia koa* pods, 5.viii.2002, PT Oboyski; Oboyski specimen PTO-129.8, genitalia slide pto-s342. 1♂ paratype (EMEC): Hawaiian Is: Maui, upper Makawao F.R., *Acacia koa* forest, 1291 m., at UV light, 3.vii.2005, PT Oboyski; Oboyski specimen PTO-529.8. 1♂ paratype (EMEC): Hawaiian Is: Maui, Haleakala N.P., Kaupo Gap, *A. koa* forest – UV light, 1680 m., 8.vii.2005, PT Oboyski; Oboyski specimen PTO-540.10.

BIOLOGY: Larvae feed within the seedpods of *Acacia koa*. Although *Pristomerus hawaiiensis* and *Euderus metallicus*, which parasitize other *Cydia* species, emerged from seedpods containing *C. anomalosa*, it was not possible to tell whether they emerged from these or *Cryptophlebia illepada*, found in the same collection of pods. Adults are attracted to ultraviolet lights.

DISTRIBUTION: Endemic to Hawaiian Islands: Kauai – Mt Kahili, Kokee State Park; Maui – Haleakala. Sympatric with *C. conspicua*, *C. acaciavora*, and *C. walsinghamii* at some locations.



REMARKS: This species is typical of non-Hawaiian *Cydia* in having an anal roll and without a sex pouch on the hindwings. It is unlikely that these characters would re-evolve after being lost and appears to represent a separate lineage of *Cydia* in Hawaii. However, mitochondrial and nuclear DNA evidence suggest this species is closely related to other *Acacia*-feeding Hawaiian *Cydia* (Oboyski Chapter 3).



Figures 1-4. Eggs, larvae, parasitoids, and pupa of *Cydia plicata* (Walsingham). 1. Section of *Sophora chrysophylla* seedpod with arrows indicating a cluster of three eggs and a single egg. 2. Larva feeding inside *S. chrysophylla* seed. 3. Larvae of parasitoid *Euderus metallicus* (Ashmead 1901) feeding on *Cydia* larva. 4. Pupal exuvia partially protruding from *S. chrysophylla* seedpod.



Figures 5-8. Host plants of Hawaiian *Cydia*. 5. *Sophora chrysophylla* seedpods on Mauna Kea, Hawaii, arrow indicating silk plug made by *C. plicata* larva. 6. *S. chrysophylla* terminal twig at Kokee, Kauai, showing split from previous *C. makai* feeding. 7. *Acacia koa* at Kokee, Kauai, showing flowers and *Uromyces koeae* rust gall. 8. *Canavalia galeata* flowers, seedpods, and stems from Punaluu, Hawaii.



Figures 9. Wing venation of *Cydia parapteryx* (Meyrick). The pouch hangs below the wing opening along a thin slit on the upper surface of the wing along the path of the obscured CuP vein.




Figure 10. Hawaiian *Cydia* wing pattern elements. A and B. *C. koiiae* n.sp. Oboyski, *C. C. plicata* (Walsingham).












Figure 11-12. Forewing tips of Hawaiian *Cydia*. 11. Forewing tip of *Cydia mauiensis* showing sinuous termen and crescent-shaped apical patch (arrow) characteristic of the *Canavaliae* group of species. 12. Forewing tip of *Cydia walsinghamii* showing the nearly linear termen and subtriangular apical patch (arrow) typical of most Hawaiian *Cydia*.




Figures 13-18. Hindwing ventral pouch and modified pecten scales. 13. Ventral view of hindwing pouch (arrows) of *Cydia plicata*, 14. Dorsal view showing opening to pouch (arrow) below cubital in *C. plicata*, 15. Base of modified pecten scales of *C. plicata* bending into pouch, 16. Base of modified pecten scales of *C. latiferreana* from California bending into pouch, 17. Tip of modified pecten scale of *C. plicata*, 18. Tip of pecten modified scale of *C. latiferreana*.



Figures 19-24. Hindwing anal roll with modified scales. 19. *C. anomalosa* showing white scales within dorsal pocket of hindwing anal roll. 20. SEM of thecae (androconial scales) along vein A3 in male hindwing of *C. latiferreana*. 21. SEM of hindwing anal roll of *C. anomalosa*. 22. SEM of hindwing anal roll of *C. latiferreana*. 23. SEM of modified scales along the margin of *C. anomalosa* anal roll showing phylliform sex scales. 24. SEM of modified scales along the margin of *C. latiferreana* anal roll showing phylliform and fusiform sex scales.



Figures 25-37. Hawaiian *Cydia* adult habitus. 25. *C. mauiensis*. 26. *C. falsifalcella*. 27. *C. falsifalcella*. 28. *C. parapteryx*. 29. *C. velocilimitata*. 30. *C. velocilimitata*. 31. *C. plicata*. 32. *C. plicata*. 33. *C. plicata*. 34. *C. haleakalaensis*. 35. *C. latifemoris*. 36. *C. makai*. 37. *C. makai*




Figures 38-51. Hawaiian *Cydia* adult habitus (continued). 38. *C. koaiae*. 39. *C. koaiae*. 40. *C. conspicua*. 41. *C. conspicua*. 42. *C. rufipennis*. 43. *C. rufipennis*. 44. *C. montana*. 45. *C. montana*. 46. *C. montana*. 47. *C. hawaiiensis*. 48. *C. acaciavora*. 49. *C. walsinghamii*. 50. *C. anomalousa*. 51. *C. anomalousa*


52



53



54




55



56




57




Figures 52-57. Male genitalia 52. *Cydia gypsograptia* – male holotype, slide BM 9543. 53. *C. mauiensis* – male holotype, slide PTO-s254. 54. *C. falsifalcella* – male holotype, slide BM 2051. 55. *C. parapteryx* – slide PTO-s252. 56. *C. velocilimitata* – slide PTO-s170. 57. *C. plicata* – male holotype, slide BM 1885.


58



59




60




61




62









63



Figures 58-63. Male genitalia (continued). 58. *Cydia haleakalaensis* – slide PTO-s339. 59. *C. latifemoris* – allotype, slide BM 2053. 60. *C. makai* – slide PTO-s311. 61. *C. rufipennis* – slide BM 2058. 62. *C. montana* – slide PTO-s323. 63. *C. koaiae* – slide PTO-s333.



Figures. 64-69. Male genitalia (continued). 64. *Cydia conspicua* – slide PTO-s335. 65. *C. hawaiiensis* – slide PTO-s302. 66. *C. acaciavora* – slide PTO-s307. 67. *C. walsinghamii* – slide BM 1892. 68. *C. crassicornis* – holotype, slide BM 1880). 69. *C. anomalousa* – slide PTO-s342.




Figures 70-78. Female genitalia. 70. *Cydia chlorostola* – holotype, slide BM 9546 Clarke. 71. *C. mauiensis* – slide PTO-s154. 72. *C. falsifalcella* – slide PTO-s183. 73. *C. parapteryx* – paralectotype, slide BM 7549. 74. *C. plicata* – allotype, slide BM 1886. 75. *C. obliqua* – holotype, slide BM 1882. 76. *C. storeella* – holotype, slide BM 1881. 77. *C. haleakalaensis* – slide PTO-s318. 78. *C. latifemoris* – holotype, slide BM 2054.


79




80




81




82




83




84



85



86



Figures 79-86. Female genitalia (continued). 79. *Cydia makai* – slide PTO-s147. 80. *C. rufipennis* – slide BM 2057. 81. *C. montana* – slide PTO-s322. 82. *C. conspicua* – slide PTO-s336. 83. *C. hawaiiensis* – slide PTO-s42. 84. *C. acaciavora* – slide PTO-s315. 85. *C. walsinghamii* – slide PTO-s308. 86. *C. anomalosa* – slide PTO-s169.

Literature Cited

- Banko, P. C., P. T. Oboyski, J. W. Slotterback, S. J. Dougill, D. M. Goltz, L. Johnson, M. E. Laut, and T. C. Murray. 2002. Availability of food resources, distribution of invasive species, and conservation of a Hawaiian bird along a gradient of elevation. *Journal of Biogeography* 29(5-6):789-808.
- Brenner, G. J., P. T. Oboyski, and P. C. Banko. 2002. Parasitism of *Cydia* spp. (Lepidoptera: Tortricidae) on *Sophora chrysophylla* (Fabaceae) along an elevation gradient of dry subalpine forest on Mauna Kea, Hawaii. *Pan-Pacific Entomologist* 78(2):101-109.
- Bridwell, J. C. 1919. Some notes on Hawaiian and other Bethyridae (Hymenoptera) with descriptions of new species. *Proceedings of the Hawaiian Entomological Society* 4:21-38.
- Butler, A. G. 1881. On a collection of nocturnal Lepidoptera from the Hawaiian Islands. *Annals and Magazine of Natural History* 5(7):317-333, 392-408.
- Butler, A. G. 1882. On a small collection of Lepidoptera from the Hawaiian Islands. *Transactions of the Entomological Society of London* 1882:31-45.
- Evenhuis, N. L., editor. 2007. Barefoot on lava: The journals and correspondence of naturalist R.C.L. Perkins in Hawaii, 1892-1901. Bishop Museum Press, Honolulu.
- Horak, M. 2006. Olethreutine moths of Australia (Lepidoptera: Tortricidae). *Monographs on Australian Lepidoptera* 10:522.
- Komai, F. 1999. A taxonomic review of the genus *Grapholita* and allied genera (Lepidoptera: Tortricidae) in the Palearctic region. *Entomologica Scandinavica (Supplement)* 55:4-226.
- Komai, F. and M. Horak. 2006. *Grapholitini* Guenée. Pages 396-467 in M. Horak, editor. *Olethreutine moths of Australia*. CSIRO Publishing, Canberra.
- Meyrick, E. 1928. Some new species of Hawaiian Lepidoptera. *Proceedings of the Hawaiian Entomological Society* 7(1):91-104.
- Meyrick, E. 1932. Exotic microlepidoptera. IV:193-352.
- Miller, W. E. 1990. Body size and diet quality in the genus *Cydia* (Tortricidae). *Journal of the Lepidopterists' Society* 44:113-142.
- Namba, R. 1957. *Cryptophlebia illepida* (Butler) (Lepidoptera: Eucosmidae) and other insect pests of the macadamia nut in Hawaii. *Proceedings of the Hawaiian Entomological Society* 16(2):284-297.
- Oboyski, P. T., J. W. Slotterback, and P. C. Banko. 2004. Differential parasitism of seed-feeding *Cydia* (Lepidoptera: Tortricidae) by native and alien wasp species relative to elevation in subalpine *Sophora* (Fabaceae) forests on Mauna Kea, Hawaii. *Journal of Insect Conservation* 8(2-3):229-240.
- Perkins, R. C. L. 1913. Introduction. Pages xv-ccxxviii in D. Sharp, editor. *Fauna Hawaiiensis*. Cambridge University Press, Cambridge.

- Pratt, T. K., C. T. Atkinson, P. C. Banko, J. D. Jacobi, and B. L. Woodworth, editors. 2009. Conservation biology of Hawaiian forest birds: Implications for island avifauna. Yale University Press, New Haven.
- Robinson, G. S. 1976. The preparation of slide of Lepidoptera genitalia with special reference to Microlepidoptera. *Entomologist's Gazette* 27:127-132.
- Stein, J. D. 1983a. The biology, host range, parasites, and hyperparasites of koa seed insects in Hawaii: a review. *Proceedings of the Hawaiian Entomological Society* 24(2/3):317-326.
- Stein, J. D. 1983b. Insects associated with *Acacia koa* seed in Hawaii. *Environmental Entomology* 12:299-302.
- Swezey, O. H. 1908. Life history notes of two variable tortricids. *Proceedings of the Hawaiian Entomological Society* 2:14-16.
- Swezey, O. H. 1936. Fruit-eating and seed-eating insects in Hawaii. *Proceedings of the Hawaiian Entomological Society* 9(2):196-202.
- Swezey, O. H. 1954. Forest Entomology in Hawaii. An annotated check-list of the insect faunas of the various components of the Hawaiian forests [B.P. Bishop Museum Special Publication 44]. Honolulu Star-Bulletin, Honolulu
- Swezey, O. H. and F. X. Williams. 1932. Some observations on forest insects at the Nauhi nursery and vicinity on Hawaii. *Proceedings of the Hawaiian Entomological Society* 8:179-192.
- Walsingham, L. 1907. Microlepidoptera. Pages 469-759 in D. Sharp, editor. *Fauna Hawaiiensis*. Cambridge University Press, Cambridge.
- Zimmerman, E. C. 1978. Insects of Hawaii, Volume 9: Microlepidoptera, Part I. University of Hawaii Press, Honolulu.